

Sensitivity estimates on anomalous couplings of the tau-lepton in pp , e^-p and e^+e^- colliders

A. Gutiérrez-Rodríguez,^{a,*} M. Köksal,^b A. A. Billur^c and M. A. Hernández-Ruíz^d

^aUnidad Académica de Física, Universidad Autónoma de Zacatecas,
Apartado Postal C-580, 98060 Zacatecas, México.

^bDepartment of Optical Engineering, Sivas Cumhuriyet University,
58140, Sivas, Turkey.

^cDepartment of Physics, Sivas Cumhuriyet University,
58140, Sivas, Turkey.

^dFacultad de Ciencias Químicas, Universidad Autónoma de Zacatecas
Apartado Postal 585, 98060 Zacatecas, Zac. México.

E-mail: alexgu@fisica.uaz.edu.mx, mahernan@uaz.edu.mx

We explore the discovery prospect of the pp , e^-p and e^+e^- colliders at the LHC, FCC-he and CLIC for model-independent sensitivity estimates on the dipole moments of the τ -lepton \tilde{a}_τ and \tilde{d}_τ . We consider the channels $pp \rightarrow p\tau^+\tau^-\gamma p$, $e^-p \rightarrow e^-\tau^+\tau^-\gamma p$ and $e^+e^- \rightarrow e^+\tau^+\tau^-\gamma e^-$. Bounds on the dipole moments of the τ -lepton are obtained: $\tilde{a}_\tau = (-0.00674, 0.00658)$, $|\tilde{d}_\tau(\text{ecm})| = 3.6925 \times 10^{-17}$ [LHC]; $\tilde{a}_\tau = (-0.00265, 0.00246)$, $|\tilde{d}_\tau(\text{ecm})| = 1.4379 \times 10^{-17}$ [FCC-he] and $\tilde{a}_\tau = (-0.00128, 0.00105)$, $|\tilde{d}_\tau(\text{ecm})| = 6.4394 \times 10^{-18}$ [CLIC]. The sensitivity of CLIC show a potential advantage compared to those from LHC and FCC-he.

41st International Conference on High Energy physics - ICHEP2022
July 6 - July 13, 2022
Bologna, Italy

*Speaker

1. Introduction

The study of the τ -lepton by the ATLAS and CMS Collaborations at the Large Hadron Collider (LHC) has developed significantly in recent years and now represents a very active physics program.

The future e^-p colliders, such as the Large Hadron Electron Collider (LHeC) [1] and the Future Circular Collider Hadron Electron (FCC-he) [1], are a hybrid between the pp and e^+e^- colliders and will complement the physics program of the LHC.

In this work, we have based our study on three phenomenological analyses for finding physics beyond the Standard Model (BSM) at present and future colliders in order to compare the electromagnetic properties of the τ -lepton. We consider pp collisions at the LHC with 13, 14 TeV and luminosities 10, 30, 50, 100, 200 fb^{-1} . Another scenario is the FCC-he with 7.07, 10 TeV and $\mathcal{L} = 100, 300, 500, 700, 1000 \text{ fb}^{-1}$. The CLIC at CERN is another option with 1.5, 3 TeV and assumed luminosities 100, 300, 500, 1000, 1500, 2000, 3000 fb^{-1} .




Fig. 1: A schematic diagram for the processes $pp(e^-p, e^+e^-) \rightarrow p\tau\bar{\tau}\gamma p(e^- \tau\bar{\tau}\gamma p, e^+ \tau\bar{\tau}\gamma e^-)$.

2. The total cross-sections in pp , e^-p and e^+e^- colliders

The most general expression for the vertex of interaction $\tau\bar{\tau}\gamma$ is given by [2–5]:

$$\Gamma_\tau^\alpha = eF_1(q^2)\gamma^\alpha + \frac{ie}{2m_\tau}F_2(q^2)\sigma^{\alpha\mu}q_\mu + \frac{e}{2m_\tau}F_3(q^2)\sigma^{\alpha\mu}q_\mu\gamma_5 + eF_4(q^2)\gamma_5(\gamma^\alpha - \frac{2q^\alpha m_\tau}{q^2}). \quad (1)$$

We evaluate the total cross-sections $\sigma(pp \rightarrow p\tau\bar{\tau}\gamma p)$, $\sigma(e^-p \rightarrow e^-\tau\bar{\tau}\gamma p)$ and $\sigma(e^+e^- \rightarrow e^+\tau\bar{\tau}\gamma e^-)$ to probe the dipole moments \tilde{a}_τ and \tilde{d}_τ , with the potential of LHC, FCC-he and CLIC based $\gamma^*\gamma^*$ colliders. Furthermore, in order to suppress the backgrounds and optimize the signal sensitivity, we impose the following kinematic basic acceptance cuts for $\tau^+\tau^-\gamma$ events at the LHC, FCC-he and CLIC:

$$\begin{aligned} p_t^\gamma &> 20 \text{ GeV}, & |\eta^\gamma| &< 2.5, \\ p_t^{\tau^+, \tau^-} &> 20 \text{ GeV}, & |\eta^{\tau^+, \tau^-}| &< 2.5, \\ \Delta R(\tau^-, \gamma) &> 0.4, & \Delta R(\tau^+, \tau^-) &> 0.4, & \Delta R(\tau^+, \gamma) &> 0.4. \end{aligned} \quad (2)$$

2.1 Cross-section of the $pp \rightarrow p\gamma^*\gamma^*p \rightarrow p\tau^+\tau^-\gamma p$ signal at the LHC

The total cross-section of the process $pp \rightarrow p\gamma^*\gamma^*p \rightarrow p\tau^+\tau^-\gamma p$ signal as a function of the anomalous parameters κ and $\tilde{\kappa}$ are given in Fig. 2.




Fig. 2: The total cross-sections of the process $pp \rightarrow p\tau\bar{\tau}\gamma p$ as a function of κ and $\tilde{\kappa}$ for center-of-mass energy of (a) $\sqrt{s} = 13$ TeV and (b) $\sqrt{s} = 14$ TeV at the LHC.

2.2 Cross-section of the $e^-p \rightarrow e^-\gamma^*\gamma^*p \rightarrow e^-\tau^+\tau^-\gamma p$ signal at FCC-he

The total cross-section of the process $e^-p \rightarrow e^-\gamma^*\gamma^*p \rightarrow e^-\tau^+\tau^-\gamma p$ as a function of the two independent anomalous couplings κ and $\tilde{\kappa}$, with 7.07 and 10 TeV at FCC-he is given in Fig. 3.




Fig. 3: The total cross-sections of the process $e^-p \rightarrow e^-\tau\bar{\tau}\gamma p$ as a function of κ and $\tilde{\kappa}$ for center-of-mass energy of (a) $\sqrt{s} = 7.07$ TeV and (b) $\sqrt{s} = 10$ TeV at the FCC-he.

2.3 Cross-section of the $e^+e^- \rightarrow e^+\gamma^*\gamma^*e^- \rightarrow e^+\tau^+\tau^-\gamma e^-$ signal at CLIC

The total cross-section for the $e^+e^- \rightarrow e^+\gamma^*\gamma^*e^- \rightarrow e^+\tau^+\tau^-\gamma e^-$ processes at CLIC is given in followings figures:




Fig. 4: The total cross-sections of the process $e^+e^- \rightarrow e^+\tau\bar{\tau}\gamma e^-$ as a function of κ and $\tilde{\kappa}$ for center-of-mass energy of (a) $\sqrt{s} = 1.5$ and (b) $\sqrt{s} = 3$ TeV at the CLIC.

3. Conclusions

With $300 fb^{-1}$ of data that will be collected by LHC, a sensitivity of $\tilde{a}_\tau = (-0.0067, 0.0065)$, 95% C.L. can be achieved for the τ -lepton, and $|\tilde{d}_\tau| = 3.692 \times 10^{-17}$ ecm, 95% C.L. can be achieved for the EDM. In the case of the FCC-he, with $1000 fb^{-1}$, it is possible to obtain a sensitivity of $\tilde{a}_\tau = (0.00265, 0.00246)$ and $|\tilde{d}_\tau| = 1.437 \times 10^{-17}$ ecm at 95% C.L.. While for the CLIC, the projections with $3000 fb^{-1}$ of data that will be collected are $\tilde{a}_\tau = (0.00128, 0.00105)$ and $|\tilde{d}_\tau| = 6.439 \times 10^{-18}$ ecm, at 95% C.L.. The precision of the τ -lepton is about 39% of the SM prediction. Thus, in this framework and with the large amount of data collected at current and future colliders, BSM can be constrained much better than before [6–9]. In summary, the future CLIC at high energy and high luminosity should provide the best sensitivity on the MDM and EDM of the τ -lepton and shows a potential advantage compared to those from LHC and FCC-he [10].

Acknowledgments

We acknowledge support of the Organizing Committee of the 41st International Conference on High Energy Physics (ICHEP 2022), Bologna, Italy.

References

- [1] J. L. Abelleira Fernandez, *et al.*, *J. Phys. G: Nucl. Part. Phys.* **G39**, 075001 (2012).
- [2] S. Eidelman and M. Passera, *Mod. Phys. Lett.* **A22**, 159 (2007).
- [3] J. A. Grifols and A. Méndez, *Phys. Lett.* **B255**, 611 (1991); Erratum *ibid.* **B259**, 512 (1991).
- [4] R. Escribano and E. Massó, *Phys. Lett.* **B395**, 369 (1997).
- [5] C. Giunti and A. Studenikin, *Phys. Atom. Nucl.* **72**, 2089 (2009).
- [6] A. Gutiérrez-Rodríguez, M. A. Hernández-Ruíz and L. N. Luis-Noriega, *Mod. Phys. Lett. A* **19** (2004), 2227
- [7] A. Gutiérrez-Rodríguez, M. A. Hernández-Ruiz and M. A. Perez, *Int. J. Mod. Phys. A* **22** (2007), 3493-3508
- [8] A. Gutiérrez-Rodríguez, *Mod. Phys. Lett. A* **25** (2010), 703-713 [arXiv:0910.4217 [hep-ph]].
- [9] A. Gutiérrez-Rodríguez, M. A. Hernández-Ruiz and C. P. Castaneda-Almanza, *J. Phys. G* **40** (2013), 035001
- [10] A. Gutiérrez-Rodríguez, M. Koksál, A. A. Billur, M. A. Hernández-Ruíz, *Int. J. Theor. Phys.* **61**, 161 (2022).