

Dark matter from sterile-sterile neutrino mixing

Pasquale Di Bari^a

*^aSchool of Physics and Astronomy, University of Southampton
Southampton, SO17 1BJ, U.K.*

E-mail: P.Di-Bari@soton.ac.uk

A solution to the problem of the origin of matter in the universe can be reasonably searched within extensions of the standard model that also explain neutrino masses and mixing. Models embedding the minimal seesaw mechanism can explain the observed matter-antimatter asymmetry of the universe via leptogenesis and dark matter via active-sterile neutrino mixing. In this case a keV lightest seesaw neutrino would play the role of warm dark matter particle. This traditional solution is now constrained by various cosmological observations. I will discuss the possibility that a much heavier but yet metastable (dark) right-handed neutrino with mass in the 1 TeV–1 PeV range can play the role of (cold) dark matter particle. The right abundance would be produced by the Higgs induced mixing with a seesaw right-handed neutrino (RHINO model), i.e., by sterile-sterile neutrino mixing. Such a mixing would necessarily require a further extension of the minimal seesaw mechanism and can be described by a dimension-five effective operator. The same mixing would also necessarily induce dark neutrino instability with lifetimes that can be much longer than the age of the universe and can escape current constraints from neutrino telescopes. Alternatively, a contribution to very high energy neutrino flux produced by dark neutrino decays could explain an anomalous excess at 100 TeV energies confirmed recently by the IceCube collaboration. I will also discuss a simple UV completion where the mediator is given by a massive fermion. Intriguingly, it comes out that the favoured scale of new physics for RHINO to satisfy the dark matter requirements coincides with the grand-unified scale: a RHINO miracle.

*Corfu Summer Institute 2022 "School and Workshops on Elementary Particle Physics and Gravity",
28 August - 1 October, 2022
Corfu, Greece*

1. Introduction

The origin of matter in the universe is one of the greatest mystery in science [1]. Our observable universe is maximally matter-antimatter asymmetric, since observations do not find evidence of primordial antimatter. The current baryonic matter abundance is then the leftover matter-antimatter asymmetry after annihilations took place in the early universe.

Combining *Planck* satellite results on CMB anisotropies and baryon acoustic oscillation data, the *Planck* collaboration finds for the baryonic contribution to the energy density parameter [2]

$$\Omega_{B0}h^2 = 0.02242 \pm 0.00014, \quad (1)$$

that translates into a total baryon-to-photon number ratio

$$\eta_{B0} \equiv \frac{n_{B0} - n_{\bar{B}0}}{n_{\gamma 0}} \simeq \frac{\Omega_{B0}, \varepsilon_{c0}}{m_p n_{\gamma 0}} \simeq 273.5 \Omega_{B0} h^2 10^{-10} = (6.12 \pm 0.04) \times 10^{-10}. \quad (2)$$

This matter-antimatter asymmetry cannot be explained within the standard model (SM) and, therefore, it is regarded as a strong motivation for new physics.

At the same time cosmological observations point to the existence of a form of matter of non-standard nature referred as dark matter (DM). Its existence is necessary at high redshifts to explain observations of galactic rotation curves, dynamics of galaxy clusters and gravitational lensing in systems like the bullet cluster. It would indeed act as a kind of invisible fluid necessary to explain gravitational interactions in these systems. At the same time, however, DM also had a crucial role primordially in order to understand structure formation and CMB temperature anisotropies. All these observations favour an interpretation of this fluid in terms of the existence of new non-standard particles. This fluid had to be cold enough¹ in order for numerical simulations to reproduce the observed large scale structure of the universe.

The cold DM contribution to the energy density parameter is today well determined both from dynamics of clusters of galaxies and from CMB anisotropies and from *Planck* + BAO data it is found [2]

$$\Omega_{DM0}h^2 = 0.11933 \pm 0.00091 \simeq 5 \Omega_{B0}h^2. \quad (3)$$

There is no SM particle with properties able to satisfy the DM requirements in order to explain the cosmological observations and for this reason the DM puzzle is also regarded as a strong motivation for the existence of new physics.

For long time the dominant solution has been represented by weakly interacting massive particles (WIMPs) with a relic abundance explained from a traditional non-relativistic freeze-out (the WIMP paradigm). This was mainly supported by the observation that the observed DM abundance could be explained for WIMP masses of order of the electroweak scale (so-called WIMP miracle). However, negative results from direct and indirect searches require a relaxation of the assumptions of the WIMP miracle paradigm, so that WIMPs, though still not excluded as possible DM candidates, do not represent any more the standard solution to the DM puzzle and many other models have been proposed and investigated. The strong constraints imposed by direct and indirect

¹This implies a free streaming (comoving) length at the matter-radiation equality time not larger than about 0.1 Mpc [3].

searches require that the new physics necessary to provide a solution to the DM puzzle must lie (i) either at energy scales higher than those accessible at the LHC or (ii) involve sufficiently small interaction couplings or (iii) some combination of the two. The solution I will discuss in this talk is indeed of this third kind since it relies both on very high energy scales and very small couplings.

The matter-antimatter asymmetry of the universe and the DM puzzles can be combined together to form the problem of the origin of matter in the universe. It is then reasonable that their solution might stem from the same kind of new physics and it is, therefore, also reasonable to investigate extensions of the SM able to address these two problems simultaneously.

From neutrino oscillations we know that neutrinos are massive and mix so that neutrino physics also provides a strong phenomenological motivation to extend the SM. It is then quite well motivated to search for a solution to the origin of matter of the universe within extensions of the SM that incorporate neutrino masses and mixing.

Here I will discuss the possibility that a right-handed (RH) neutrino in the TeV-PeV mass range can play the role of DM particle. This window is not typically considered but it is very interesting, since it is currently tested by IceCube, and more generally by neutrino telescopes.

2. Minimal type-I seesaw mechanism

Adding N RH neutrinos to the SM Lagrangian and a right-right Majorana mass term violating lepton number, one can add to the SM Lagrangian the terms ($\alpha = e, \mu, \tau; I = 1, 2, \dots, N$)

$$-\mathcal{L}_{Y+M}^{\nu} = \overline{L}_{\alpha} h_{\alpha I}^{\nu} \nu_{RI} \tilde{\Phi} + \frac{1}{2} \overline{\nu_{RI}^c} M_I \nu_{RI} + \text{h.c.}, \quad (4)$$

written in a basis where charged lepton and Majorana mass matrices are diagonal. After electroweak spontaneous symmetry breaking, we have that the neutrino mass term can be written as

$$-\mathcal{L}_m^{\nu} = \frac{1}{2} \left[(\overline{\nu}_L, \overline{\nu}_R^c) \begin{pmatrix} 0 & m_D \\ m_D^T & M \end{pmatrix} \begin{pmatrix} \nu_L^c \\ \nu_R \end{pmatrix} \right] + \text{h.c.}. \quad (5)$$

In the seesaw limit, for $M \gg m_D$, the spectrum of neutrino masses splits into a light set (ordinary neutrinos) with masses given by the seesaw formula [4]

$$\text{diag}(m_1, m_2, m_3) = U_L^{\nu\dagger} m_D \frac{1}{M} m_D^T U_L^{\nu*}, \quad (6)$$

where in the particular flavour basis we are taking the diagonalising unitary matrix U can be identified with the leptonic mixing matrix, and into a heavy set (seesaw neutrinos) with masses $M_1 \leq M_2 \leq \dots \leq M_N$. For definiteness I will consider the case $N = 3$.

If one assumes that the reheat temperature of the universe T_{RH} is not much lighter than the lightest heavy neutrino state N_1 with mass M_1 , then the N_1 decays, and possibly also those of the other heavier seesaw neutrinos, can generate a lepton asymmetry that can be partly converted into a baryon asymmetry by sphalerons if $T_{\text{RH}} \gtrsim 132 \text{ GeV}$ [5]. One has also to take into account that part of this asymmetry will be washed-out by inverse processes but it can be shown that the measured baryon-to-photon number ratio in Eq. (2) can be nicely reproduced: this is the essence of thermal leptogenesis [6], a model of baryogenesis that is a straightforward cosmological application of the seesaw mechanism [7].

It is clearly intriguing to think whether such a simple extension of the SM might also address the DM puzzle. If one writes the light and heavy mass eigenfields, that are Majorana fields, as a linear combination of left-handed (LH) and RH neutrinos, then light neutrinos are dominantly LH neutrinos and heavy neutrinos are dominantly RH neutrinos. However, they also have subdominant RH (sterile) and LH (active) components mixing together and the mixing is described by $m_D M^{-1}$. If one of the RH neutrinos has sufficiently long lifetime and at the same time be produced by such active-sterile neutrino mixing, then it could play the role of DM particle. There is a very attractive solution to such a simultaneous request [8]. If $M_1 \ll m_e$, then the N_1 lifetime is given by

$$\tau_1 \simeq 5 \times 10^{28} \text{ s} \left(\frac{M_1}{\text{keV}} \right)^{-5} \left(\frac{10^{-4}}{\theta} \right)^2, \quad (7)$$

where $\theta \equiv \sum_{\alpha} |m_{D\alpha 1}/M_1|^2$ is an effective active-sterile neutrino mixing angle. One can see that N_1 can be long enough not only to satisfy the DM long-life condition but also to evade X-ray constraints, since it would subdominantly radiatively decay. At the same time such a $\sim \text{keV}$ seesaw neutrino would mainly mix with the lightest ordinary neutrino mass eigenstate, one has specifically:

$$\nu_1 = \left[U_{1\alpha}^{\dagger} \nu_{L\alpha} + (U_{1\alpha}^{\dagger} \nu_{L\alpha})^c \right] - \left[V_{L1\alpha}^{\nu} \xi_{\alpha 1}^{\star} \nu_{R1} + (V_{L1\alpha}^{\nu} \xi_{\alpha 1}^{\star} \nu_{R1})^c \right] \quad (8)$$

$$N_1 = \left[\nu_{R1} + \nu_{R1}^c \right] + \left[\xi_{1\alpha}^T \nu_{L\alpha} + (\xi_{1\alpha}^T \nu_{L\alpha})^c \right]. \quad (9)$$

It is this mixing that is actually responsible for the N_1 -decays. At the same time it would also be responsible for the production of a N_1 -abundance given by

$$\Omega_{N_1} h^2 \sim 0.1 \left(\frac{\theta}{10^{-4}} \right)^2 \left(\frac{M_1}{\text{keV}} \right)^2, \quad (10)$$

such that the measured DM abundance in Eq. (3) can be reproduced for $\theta \sim 10^{-4}$ and $M_1 \sim \text{keV}$ [9]. Interestingly, for keV masses, the lightest seesaw neutrino would behave as warm DM, implying that compared to cold DM there would be a reduced power in the large scale structure at scales corresponding to dwarf galaxies ($\sim 0.1 \text{ Mpc}$ in comoving length). This would also help in solving some potential issues in pure cold DM N-body simulations that seem to predict too many satellite galaxies, in a galaxy like ours, compare to what is observed astronomically. At the same time warm DM would also smooth the cusp profile in galaxies that is predicted by cold DM N-body simulations but that is in tension with different observations.

On the other hand, X-ray observations place an upper bound to the mass M_1 , while galaxy distribution constraints require that the N_1 's are not too warm DM particles and place a lower bound on M_1 . These constraints have progressively closed down the allowed window ruling out such a minimal solution. However, if a large pre-existing lepton asymmetry, $L \sim 10^{-5}-10^{-4}$, is present prior to the mixing, then the DM production is resonantly enhanced [10, 11] and this can reconcile the DM requirements with the X-ray observations. Moreover, it is very interesting that the observation of a new 3.5 keV line in the X-ray observations of different clusters of galaxies [12, 13] could be explained by such a mechanism for a decaying 7 keV sterile neutrino with a mixing angle $\theta \simeq 4 \times 10^{-6}$ and its relic abundance can explain the observed DM density parameter for a lepton asymmetry at the resonance $L \simeq 4.6 \times 10^{-4}$ [14]. However, authors of more recent observations of the 3.5 keV line claimed to exclude an interpretation in terms of DM decays though this seems

to be currently controversially debated [15]. It is fair to conclude that more observations will be likely necessary to definitively test this exciting possibility, in particular the XRISM satellite should provide in the next years a final answer to the 3.5 keV anomaly [16].

Finally, let us comment that such a solution to the DM puzzle can also be combined with leptogenesis from the mixing of two heaviest seesaw neutrinos (N_2 and N_3 in our notation) with GeV masses [17, 18] that would be also responsible for the atmospheric and solar neutrino mass scales thanks to the seesaw formul. In this way one would realise a unified picture of neutrino masses, DM and leptogenesis, the so-called ν MSM model. However, such unified solution seems to rely on a very strong mass degeneracy of N_2 and N_3 , at the level of $(M_3 - M_2)/(M_2 + M_3) \simeq 10^{-16}$ [19]. The FASER experiment should be able to test the existence of GeV RH neutrinos during next years. These should be produced in B meson decays for masses $M_{2,3} \lesssim M_B \sim 3$ GeV [20].

3. Heavy RH neutrino as DM

For very heavy DM particles ($M_{\text{DM}} \gtrsim 1$ TeV), just a tiny non-thermal abundance is sufficient to reproduce the observed energy density parameter, since one has:

$$N_{\text{DM}} \sim 10^{-9} (\Omega_{\text{DM}0} h^2) N_{\gamma}^{\text{prod}} \frac{\text{TeV}}{M_{\text{DM}}} \sim 10^{-10} N_{\gamma}^{\text{prod}} \frac{\text{TeV}}{M_{\text{DM}}}. \quad (11)$$

This simple observation is encouraging, since it seems that very small couplings would be sufficient to produce such an abundance and this should make easier to have a metastable DM particle. If one imposes that one of the seesaw neutrino in the model (4) is the DM particle, then necessarily the neutrino Yukawa matrix must be in one of these three forms [21, 22]

$$h^{\nu} \simeq \begin{pmatrix} \varepsilon_{e1} & h_{e2} & h_{e3} \\ \varepsilon_{\mu1} & h_{\mu2} & h_{\mu3} \\ \varepsilon_{\tau1} & h_{\tau2} & h_{\tau3} \end{pmatrix}, \text{ or } \begin{pmatrix} h_{e1} & \varepsilon_{e2} & h_{e3} \\ h_{\mu1} & \varepsilon_{\mu2} & h_{\mu3} \\ h_{\tau1} & \varepsilon_{\tau2} & h_{\tau3} \end{pmatrix}, \text{ or } \begin{pmatrix} h_{e1} & h_{e2} & \varepsilon_{e3} \\ h_{\mu1} & h_{\mu2} & \varepsilon_{\mu3} \\ h_{\tau1} & h_{\tau2} & \varepsilon_{\tau3} \end{pmatrix}, \quad (12)$$

where the Yukawa couplings $\varepsilon_{\alpha I}$ of one RH neutrino, corresponding to the entries of one of the three columns, are tiny, possibly because proportional to some small symmetry breaking parameters. We will refer to such very weakly coupled RH neutrino as the *dark neutrino* and denote it by N_{D} . The lifetime of the dark neutrino is in this case given by

$$\tau_{\text{D}} = \frac{4\pi}{h_{\text{A}}^2 M_{\text{D}}} \sim 10^{-26} h_{\text{A}}^2 \frac{\text{TeV}}{M_{\text{D}}} \text{ s}, \quad (13)$$

where $h_{\text{A}}^2 = \sum_{\alpha} \varepsilon_{\alpha \text{D}}^2$ and the subscript D = 1, 2 or 3, depending on the choice in (12). Imposing the lower bound $\tau_{\text{D}} \gtrsim \tau_{\text{D}}^{\text{min}} \sim 10^{28}$ s from the IceCube neutrino telescope, as we will discuss, one finds $h_{\text{A}} \lesssim 10^{-27} \sqrt{\text{TeV}/M_{\text{D}}}$. Such small values for the Yukawa couplings make impossible to find any efficient production mechanism for the dark neutrino.²

We can then conclude that with a minimal type-I seesaw extension of the SM, one cannot find a solution for a very heavy dark neutrino playing the role of DM. We have then to consider a further extension.

²Here I am assuming $M_{\text{D}} \gg v \sim 100$ GeV. On the other hand, we have seen that if $M_{\text{D}} \ll m_e$ then the lifetime would be much suppressed compared to Eq. (13) and one finds the solution $M_{\text{D}} \sim \text{keV}$ with N_{D} produced by active-sterile neutrino mixing. In [23] the authors find that a MeV dark neutrino could satisfy the DM requirements, being efficiently produced by gauge boson decays, though most recent constraints X-ray observations seem to almost rule out this scenario.

4. Anisimov operators

The idea is to consider an effective theory where the new physics does not need to be specified but it is encoded in 5-dim (Anisimov) operators involving only RH neutrino and Higgs field [21, 24]. Therefore, one has now an effective Lagrangian given by

$$\mathcal{L}_{\text{eff}} = \mathcal{L}_{\text{SM}} + \mathcal{L}_{Y+M}^{\nu} + \mathcal{L}_A, \quad (14)$$

where

$$\mathcal{L}_A = \sum_{I,J} \frac{\lambda_{IJ}}{\Lambda} \Phi^\dagger \Phi \overline{N}_I^c N_J = \frac{\lambda_{\text{DS}}}{\Lambda} \Phi^\dagger \Phi \overline{N}_D^c N_S + \frac{\lambda_{\text{SS}}}{\Lambda} \Phi^\dagger \Phi \overline{N}_S^c N_S + \frac{\lambda_{\text{DD}}}{\Lambda} \Phi^\dagger \Phi \overline{N}_D^c N_D. \quad (15)$$

Here I am denoting by N_S one of the two seesaw neutrinos that reproduce the atmospheric and solar neutrino mass scales. We can just consider the interaction with one of them, since the inclusion of the third one would just produce more stringent constraints. However this still has a cosmological role in producing the necessary interference with N_S to have non-vanishing CP asymmetries. In this way the decays of the two seesaw neutrinos can produce a $B - L$ asymmetry that is partly converted by sphalerons into the observed matter-antimatter asymmetry, so that one can also have successful leptogenesis.

The first term in \mathcal{L}_A is the RH-RH (sterile-sterile) Higgs induced neutrino mixing (RHINO) operator and will be responsible for the direct production of the dark neutrinos. Before discussing this production three interesting things can be noticed:

- The effective Lagrangian can be regarded as a step further compared to SMEFT [25], where also RH neutrino fields are included in the effective operators and it can be regarded as a specific example of ν SMEFT [26].
- The Anisimov operators are analogous to the Weinberg operator, a kind of further step in energy scale.
- They are a generalisation of the usual Higgs portal renormalisable operator involving a scalar [27].

5. RHINO dark matter

I want now to focus on the RHINO operator, showing how its presence in the early universe can convert a small fraction of source neutrinos into dark neutrinos. One has then to check whether the dark neutrino relic abundance can match the observed DM abundance. First of all it is convenient to introduce the effective scale $\tilde{\Lambda}_{\text{DS}} \equiv \Lambda/\lambda_{\text{DS}}$. If $\lambda_{\text{DS}} \neq 0$, then a source-dark (sterile-sterile) neutrino mixing is induced. One has to take into account finite temperature medium effects induced by the RHINO operator. These generate a contribution to the self-energy of the mixed source-dark neutrino system (see left panel in Fig. 1). Also source neutrino Yukawa self-interactions contribute to the self energy and need to be taken into account (see right panel in Fig. 1). These self-energy contributions




Figure 1: Self-energy diagrams from Yukawa interactions (panel (a)) and from Anisimov interactions (panels (b) and (c)) [28].

can be also conveniently described in terms of effective potentials V_{DS}^Λ and V_{SS}^Y , respectively, given by

$$V_{\text{DS}}^\Lambda = \frac{T^2}{12 \tilde{\Lambda}_{\text{DS}}} \quad \text{and} \quad V_{\text{SS}}^Y = \frac{T^2}{8 E_J} h_S^2. \quad (16)$$

These need to be taken into account in the effective mixing Hamiltonian describing the evolution of the mixed system given by

$$\Delta \mathcal{H} \simeq \begin{pmatrix} -\frac{\Delta M^2}{4p} - \frac{T^2}{16p} h_S^2 & \frac{T^2}{12 \tilde{\Lambda}_{\text{DS}}} \\ \frac{T^2}{12 \tilde{\Lambda}_{\text{DS}}} & \frac{\Delta M^2}{4p} + \frac{T^2}{16p} h_S^2 \end{pmatrix}, \quad (17)$$

where we used the ultrarelativistic approximation³ and defined $\Delta M^2 \equiv M_S^2 - M_D^2$.

The production of dark neutrinos can be described by the density matrix equation [29]

$$\frac{d\mathcal{N}}{dz} = -\frac{i}{H(z)z} [\Delta \mathcal{H}, \mathcal{N}] - \begin{pmatrix} 0 & \frac{1}{2}(D+S) \mathcal{N}_{\text{DS}} \\ \frac{1}{2}(D+S) \mathcal{N}_{\text{SD}} & (D+S)(\mathcal{N}_{N_S} - \mathcal{N}_{N_S}^{\text{eq}}) \end{pmatrix}, \quad (18)$$

where we defined $z \equiv M_D/T$ normalised the density matrix \mathcal{N} in a way that the diagonal elements give the abundances of dark neutrinos, $N_{N_D} = \mathcal{N}_{\text{DD}}$, and source neutrinos, $N_{N_S} = \mathcal{N}_{\text{SS}}$. Notice

³This assumes that the dark neutrino production occurs in the ultra-relativistic regime, as it will be verified.




Figure 2: Evolution of the DM abundance $N_{N_{DM}}$ for three different choices of T_{RH} and τ_{DM} as indicated in a linear plot for the abundance (from [29]).

how decays and scatterings also contribute to decoherence effects, damping the density matrix off-diagonal terms. Initially, for $z = z_{in}$, the density matrix is simply given by:

$$\mathcal{N}(z_{in}) = N_{N_S}(z_{in}) \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}. \quad (19)$$

Here notice that the abundances are normalised in a way that in ultrarelativistic thermal equilibrium one has $N_{N_i}^{eq}(T \gg M_i) = 1$ and consequently $N_\gamma = 4/3$. Notice that with this normalisation the final value of dark neutrino abundance that is necessary to reproduce the measured value of DM energy density parameter is given by

$$N_{N_D}^{f,obs} \simeq 1.1 \times 10^{-7} \frac{\text{GeV}}{M_D}. \quad (20)$$

In Fig. 2 we show the production of the dark neutrino abundance for fixed values $M_D = 220 \text{ TeV}$ and $M_S = 300 \text{ GeV}$ and for three choices of the values of the parameters T_{RH} and τ_D as indicated, such that the relic abundance reproduces the value $N_{N_D}^{f,obs} \simeq 5 \times 10^{-13}$. Moreover, as we will see, the values of the lifetime τ_D respect the lower bound from IceCube measurement of the high energy neutrino flux. There are two important things to notice in these solutions:

- An *initial thermal abundance of source neutrinos*, i.e., $N_{N_S}^{in} = 1$, is assumed. I will show soon how this assumption can be justified within a logical extension of the scenario.
- The dark neutrino is assumed to be heavier than the source neutrino, i.e., $M_D > M_S$. All results I present in this talk are obtained for such a choice. The case $M_D < M_S$ requires a dedicated analysis that will be presented elsewhere.




Figure 3: Diagrams for two (left) and four (right) body decay (from [29]).

6. Constraints from decays

The RHINO operator is also responsible for the instability of the dark neutrinos that have necessarily to decay [21, 22]. With the assumption of heavy RH neutrinos, $M_I \gtrsim 100$ GeV, one has two dominant decay channels. The first is *two body decay* channel where the dark neutrino, via the mixing with the source neutrino, decays into a gauge boson $A = Z_0, W^\pm, \gamma, H$ and a charged lepton or neutrino with decay rate $\Gamma_{D \rightarrow A + \ell_S}$. The second channel is *four body decay*, such that $N_D \rightarrow 3A + N_S$, with decay rate $\Gamma_{D \rightarrow 3A + \ell_S}$. The diagrams for these two channels are shown in the left and right panel, respectively, of Fig. 3. Three body decays are also possible but give a sub-dominant channel that can be neglected.

In the case of the two body decay, the dark neutrino mixes today, at zero temperature, with the source neutrino with a mixing angle

$$\theta_{\Lambda 0} = \frac{2v^2 / \tilde{\Lambda}_{DS}}{M_D (1 - M_S / M_D)}. \quad (21)$$

In this way the two body decay rate is given by

$$\Gamma_{D \rightarrow A + \ell_S} = \frac{h_S^2}{\pi} \left(\frac{v^2}{\tilde{\Lambda}} \right)^2 \frac{M_D}{(M_D - M_S)^2}. \quad (22)$$

The four body decay rate can be calculated in the narrow width approximation, obtaining

$$\Gamma_{D \rightarrow 3A + \ell_S} = \frac{\Gamma_S}{15 \cdot 2^{11} \cdot \pi^4} \frac{M_D}{M_S} \left(\frac{M_D}{\tilde{\Lambda}_{DS}} \right)^2, \quad (23)$$

where $\Gamma_S = h_S^2 M_S / (4\pi)$. The lifetime can then be calculated as

$$\tau_D \simeq (\Gamma_{D \rightarrow A + \ell_S} + \Gamma_{D \rightarrow 3A + \ell_S})^{-1}. \quad (24)$$

As I am going to discuss in more detail, IceCube data impose roughly a lower bound on τ_D . As one can see from the expressions of the rates, this implies that two body decays place a lower bound on M_D while four body decay place an upper bound. In this way there is an intrinsic allowed range for the dark neutrino mass that emerges from the model when the lower bound is imposed. In Fig. 4




Figure 4: Allowed regions in the plane τ_{DM} versus M_{DM} for different values of M_S and for $T_{\text{RH}} = 10^{15}$ GeV (from [29]).

we show the allowed regions in the plane τ_{D} vs. M_{D} . Notice that there are four regions obtained for initial thermal N_S abundance. They correspond to four different values of M_S as indicated. One can see that for an increasing value of M_S , since the lifetime decreases, as it can be easily inferred from Eqs. (22)-(24), the allowed region shrinks. I will comment soon on the importance of the value of M_S for leptogenesis. First, however, at this point I notice that one could legitimately have an objection: the allowed regions have been obtained assuming arbitrarily that the source neutrino have a thermal abundance at the time of the mixing with the dark neutrinos. If we relax this assumption and consider an initially vanishing source neutrino abundance, then one can see that Yukawa interactions are by far insufficient to produce a source neutrino abundance close to a thermal value prior to the mixing. The consequence is quite dramatic: all the shown allowed regions simply evaporate. The only possibility is then to consider a value $M_S < M_W \simeq 80$ GeV in a way that the decay rate of source neutrinos decreases significantly and an allowed region appears. In Fig. 4 the case $M_S = 1$ GeV is shown. One can see that the allowed region now appears for $M_{\text{D}} \gtrsim 10$ PeV, so for much heavier dark neutrinos. This is because the upper bound on M_{D} from four body decays essentially disappears and one is left only with a much more stringent lower bound from two body decays. The situation is clearly much less attractive now and so one can have a legitimate question: is there any process that can thermalise the source neutrino abundance prior to the mixing with the dark neutrinos thus opening the (much more attractive) allowed regions at much higher values of M_S and lower values of M_{D} ?

There are two very good motivations to try to investigate this possibility. I am going first to discuss these two motivations and then as a last point of my talk I will finally show that indeed there is a solution to such an issue and it is actually quite a natural one, not artificially imposed.




Figure 5: Evolution of the $B - L$ asymmetry and DM abundance $N_{N_{DM}}$ for a choice of parameters such that the final values simultaneously reproduce the observed values of matter-antimatter asymmetry and DM abundance. The N_S -abundance is also shown. One can see that the plot is for initial thermal N_S -abundance (from [29]).

7. Unified picture for the origin of matter in the universe

The value of M_S is particularly interesting if one wants to combine leptogenesis, solving the problem of the origin of matter within the same model, since it sets the scale of leptogenesis. It can be seen how for value above 10 TeV the allowed regions in the case of initially thermal N_S abundance tend to disappear. The minimum value of M_S for (resonant) leptogenesis to work in a way to be independent of the initial conditions is approximately $M_S \sim 300$ GeV, since below this value, the $B - L$ asymmetry is generated in the non-relativistic regime, at the end of the wash-out regime, too late to be converted into a baryon asymmetry. One has in this case to resort to contributions produced in the ultra-relativistic regime that strongly depend on the initial conditions. Therefore, there is a wide allowed range $M_S \sim 300$ GeV–100 TeV that allows a solution to the problem of the origin of matter of the universe if one can justify an initial thermal N_S abundance. In Fig. 5 one can see a particular realisation of a simultaneous generation of dark neutrino abundance and $B - L$ asymmetry that reproduce the measured values of DM energy density parameter and baryon-to-photon number ratio. The possibility to realise such a picture is particularly intriguing but one needs to find a way to thermalise the source neutrinos prior to the mixing. Let us first discuss a second equally intriguing motivation that is related to the possibility to test the RHINO model.

8. IceCube data and the 100 TeV anomaly

Neutrinos are perfect astronomical messengers since they can travel to us from the edge of the universe [30]. In the energy range 10 TeV–10 EeV they are the only particles that can travel unabsorbed and undeflected. In 2013 IceCube discovered very high energy neutrinos (30 TeV – 1 PeV range) [31]. Some have been observed in coincidence with blazar γ -ray flare, an evidence of the presence of a component with extragalactic origin. IceCube employs two strategies to reduce the overwhelming atmospheric background at energies $\lesssim 300$ TeV. A first strategy is to impose a High Energy Starting Events (HESE) veto. This sample is the first that found evidence of a diffuse extraterrestrial neutrino component. A second strategy, more traditional, is based on analysing an up-going muon data set. This has now also confirmed the existence of a diffuse extraterrestrial neutrino component. However, there is a tension between the two data sets. The up-going muon sample energy spectrum is well described by a single power law $\propto E^{-\gamma_{\text{astro}}}$ with spectral index $\gamma_{\text{astro}} = -2.19$ [32]. This is quite a standard astrophysical component well explained by a Fermi model for the acceleration of cosmic rays. On the other hand, the case of HESE data, one would need a two-component power law spectrum, one *hard* component with a value of the spectral index compatible with the one from up-going muon sample dominating at energies above ~ 300 TeV and a second *soft* component with spectral index $\gamma_{\text{soft}} \simeq 3.7$ dominating at energies $E \sim 100$ TeV [33]. This second component is more difficult to understand in astrophysical models but not particularly surprising. What makes it challenging to be understood in terms of an astrophysical solution is that such a soft component would be incompatible within a multimessenger analysis showing that the γ -ray flux at ~ 100 GeV energies (as measured by the Fermi satellite) would be comparable with the hard component flux as expected within traditional astrophysical models but it would be incompatible with the soft component that seems to suggest the existence of *hidden* sources producing just neutrinos and not photons. Similar conclusion can be also drawn analysing ultra-high energy cosmic rays spectrum. Unknown and quite exotic astrophysical hidden sources might be a solution to this anomaly. It is also intriguing that such unexplained excess can be well reproduced within the RHINO model [22]. In Fig. 6 one can see an example how the decays of dark neutrino with mass $M_D = 300$ TeV can reproduce the excess and help fitting IceCube data (in this 2016 analysis, ~ 4 year HESE data were employed). Recently, the IceCube collaboration has presented an analysis where IceCube 7.5 year data are fitted including a contribution to the neutrino flux from decaying DM into various channels, finding that with such an addition the fit improves at 2.5σ significance level compared to the null hypothesis, corresponding to the case where just a power law astrophysical component is present [34]. A decaying DM scenario it is then a viable way to explain this excess at 100 TeV. Moreover, collection of more data should allow to perform tests on the level of the anisotropies in the flux of very high energy neutrinos. This would allow to disentangle astrophysical explanations in terms of exotic hidden sources from a DM scenario, since anisotropies should show the presence of a component tracking the DM distribution.

In conclusion, the possibility to combine leptogenesis with RHINO DM in a unified model of the origin of matter of the universe and the possibility to provide an explanation for the 100 TeV excess in very high energy neutrino data, provide two strong motivations to answer the question whether it is possible to find processes that can thermalise the source neutrino abundance prior to the dark neutrino production from their mixing.




Figure 6: Event spectra in the IceCube detector after 1347 days including a contribution from RHINO DM decays with $M_D = 300$ TeV (from [22]).

9. Including Higgs portal interactions for the source neutrino: a RHINO miracle

Let us go back to the full set of Anisimov operators in Eq. (15). We have so far neglected Higgs portal interactions producing the source neutrinos. These are natural processes to be included that might lead to a thermalisation of the source neutrinos prior to the mixing with the dark neutrinos [28]. Therefore, we now consider the effective Lagrangian:

$$\mathcal{L}_{\text{eff}} = \mathcal{L}_{\text{SM}} + \mathcal{L}_{Y+M}^\nu + \frac{1}{\tilde{\Lambda}_{\text{DS}}} \Phi^\dagger \Phi \bar{N}_D^c N_S + \frac{1}{\tilde{\Lambda}_{\text{SS}}} \Phi^\dagger \Phi \bar{N}_S^c N_S, \quad (25)$$

where $\tilde{\Lambda}_{\text{SS}} \equiv \lambda_{\text{SS}}/\Lambda$. The density matrix equation (26) now needs to be modified to include these additional interactions, becoming:

$$\frac{d\mathcal{N}}{dz} = -\frac{i}{H(z)z} [\Delta\mathcal{H}, \mathcal{N}] - \begin{pmatrix} 0 & \frac{1}{2}(D+S) \mathcal{N}_{\text{DS}} \\ \frac{1}{2}(D+S) \mathcal{N}_{\text{SD}} & (D+S)(N_{N_S} - N_{N_S}^{\text{eq}}) + A(N_{N_S}^2 - N_{N_S}^{\text{eq}2}) \end{pmatrix}, \quad (26)$$

where

$$A \equiv \frac{\langle \sigma_{\phi\phi^\dagger \rightarrow N_S N_S^c} v_{\text{rel}} \rangle}{H(z)z R^3(z)}, \quad (27)$$

and $\langle \sigma_{\phi\phi^\dagger \rightarrow N_S N_S^c} v_{\text{rel}} \rangle$ is the thermal averaged cross section. Here $R^3(z)$ is the portion of comoving volume where abundances are calculated, essentially a normalisation factor.

The thermal averaged cross section is simply given by [35]

$$\langle \sigma_{\phi\phi^\dagger \rightarrow N_S N_S^c} v_{\text{rel}} \rangle \Big|_{M_S \ll T} = \frac{1}{4\pi \tilde{\Lambda}_{\text{SS}}^2}. \quad (28)$$




Figure 7: Effect of source neutrino Higgs portal interactions on the source and dark neutrino abundances (from [28]).

One can rewrite $A(z)$ as

$$A(z) = \frac{A_1}{z^2}, \quad \text{with } A_1 \equiv A(z=1) = \frac{3}{16} \frac{\zeta(3)}{\pi^3} g_{Ns} \sqrt{\frac{90}{8\pi^3 g_R}} \frac{M_D M_P}{\tilde{\Lambda}_{SS}^2} \quad (29)$$

and a convenient numerical expression for A_1 is given by

$$A_1 \simeq 1.0 \times 10^{-11} \left(\frac{M_D}{100 \text{ TeV}} \right) \left(\frac{10^{16} \text{ GeV}}{\tilde{\Lambda}_{SS}} \right)^2. \quad (30)$$

From this expression for $A(z)$ one can then finally calculate the source neutrino abundance prior to the mixing given by

$$N_{Ns}(z_{in} \ll z \ll 1) - N_{Ns}(z_{in}) \simeq \frac{A_1}{z_{in}} \simeq 1.0 \times \left(\frac{T_{in}}{10^{16} \text{ GeV}} \right) \left(\frac{10^{16} \text{ GeV}}{\tilde{\Lambda}_{SS}} \right)^2. \quad (31)$$

This numerical expression already highlights the emergence of the grand-unified scale as the natural scale of new physics for the effectiveness of these interactions in thermalising the source neutrinos.

In Fig. 7 we show a benchmark case of evolution of the abundances of source and dark neutrinos including both the RHINO operator and Higgs portal interactions for the source neutrinos. In this example we have taken $\tilde{\Lambda}_{SS} = 10^{16} \text{ GeV}$ and $\tilde{\Lambda}_{DS} = 10^{23} \text{ GeV}$. These are typical values leading to successful dark neutrino production and stability.

In Fig. 8 the allowed regions in the traditional plane τ_D versus M_D are shown as in Fig. 5. This time there is no assumption of initial thermal source neutrino abundance, the job is done by the Higgs portal interactions. In this figure the effective scales $\tilde{\Lambda}_{SS} = 10^{16} \text{ GeV}$ and $M_S = 300 \text{ GeV}$. The regions are shown for different value of T_{RH} . In Fig. 9 the same regions are shown for $M_S = 10 \text{ TeV}$ and $M_S = 100 \text{ TeV}$ and again one can see that the latter gives the maximum value of the seesaw scale corresponding to the scale of leptogenesis. At these scales leptogenesis needs to be resonant [36] but the final asymmetry can be independent of the initial conditions. In the figures one can also see lower bounds on τ_D placed by different data sets and assuming different decay channels




Figure 8: Allowed regions in the lifetime versus mass plane for a fixed value $\tilde{\Lambda}_{SS} = 10^{16}$ GeV, for the indicated values of T_{RH} and for $M_S = 300$ GeV (upper panel), 1 TeV (bottom panel) [28].

for DM. The star denotes the best fit found in [34] for 7.5 year IceCube HESE data. The model can easily reproduce the best fit value for $M_D = 386$ GeV. The choice of the grandunified scale for $\tilde{\Lambda}_{SS}$ maximises the allowed region in the parameter space and emerges as the favourite scale of new physics in the RHINO model and this can legitimately be seen as a very encouraging coincidence, what I refer to as the *RHINO miracle*, paraphrasing the old WIMP miracle. However, there is another important objection and potentially unsatisfactory feature. In Fig. 7 we have seen that the typical necessary values of the effective scales $\tilde{\Lambda}_{SS} \sim 10^{16}$ GeV and $\tilde{\Lambda}_{DS} \sim 10^{23}$ GeV differ by many orders of magnitudes. It would seem that a successful production of dark neutrinos able to respect all DM experimental requirements necessarily requires two different new physics scales. This complications, and increase of parameters of the model, looks as a price to pay for it to work, providing a successful picture of the origin of matter of the universe. However, as I am going to discuss, there is a simple UV-completion clearly showing it is possible to have just one scale of new physics and strikingly this coincides with the grandunified scale.

10. UV-completions: a possible GUT origin

I will now discuss two possible UV-completions for the RHINO models including Higgs portal interactions for the source neutrinos. In a first case the mediator of the interactions is a very heavy Higgs scalar [21, 22, 28, 35] and in the second case a very heavy fermion [21, 28]. As we will see, while the first option leads to quite a rather contrived choice of parameters, the second option is strikingly simple and successful in addressing all issues.

10.1 Heavy scalar as a mediator

In a first extension of the seesaw Lagrangian one introduces a heavy real scalar field H (with vanishing vev) coupling to the RH neutrinos with Yukawa couplings y_{IJ} and to the standard Higgs field with a trilinear coupling μ :

$$\mathcal{L}_H = \frac{1}{2} \partial_\mu H \partial^\mu H - \frac{1}{2} M_H^2 H^2 - \lambda_{IJ} H \overline{N}_I^c N_J - \mu H \phi^\dagger \phi. \quad (32)$$




Figure 9: Feynman diagrams with a heavy scalar H as mediator and $I, J = D, S$. Integrating out H , they lead to the Feynman diagrams in panel (b) of Fig. 2 and (a), (b) and (c) of Fig. 3.

At scales much below M_H we can integrate out H , obtaining the effective Lagrangian

$$\mathcal{L}_H^{\text{eff}} = \frac{1}{2} \sum_{I,J,K,L} \frac{\lambda_{IJ}\lambda_{KL}}{M_H^2} (\overline{N}_I^c N_J) (\overline{N}_K^c N_L) + \frac{1}{2} \frac{\mu^2}{M_H^2} (\phi^\dagger \phi)^2 + \frac{\mu \lambda_{IJ}}{M_H^2} \Phi^\dagger \Phi \overline{N}_I^c N_J. \quad (33)$$

One can clearly recognise the Anisimov operators in Eq. (15) and the effective scales can be identified with $\tilde{\Lambda}_{IJ} = \Lambda/\lambda_{IJ}$, and $\Lambda = M_H^2/\mu$. Diagrammatically, the self-energy diagram in the panel (b) of Fig. 1 and a four-point interaction scattering diagram (not shown) are obtained by the diagrams in Fig. 9, panel (a) and panel (b), respectively. The appealing feature of this model is that one can get a trans-Planckian value for the effective scale $\tilde{\Lambda}_{DS} \sim 10^{23}$ GeV even for $\lambda_{IJ} = \mathcal{O}(1)$, simply choosing $\mu \ll M_{\text{GUT}}$. For example, one can take $M_H \sim M_{\text{GUT}} \sim 10^{16}$ GeV and $\mu \sim 10^9$ GeV. However, the problem of this setup is that one cannot also reproduce the effective scale $\tilde{\Lambda}_{SS} \sim 10^{16}$ GeV for the source neutrino Higgs portal interactions. In that respect, one should arbitrarily assume $\Lambda \sim 10^{16}$ GeV, for example for $\mu = M_H \sim M_{\text{GUT}} \sim 10^{16}$ GeV, $\lambda_{DS} \sim 10^{-7}$ and $\lambda_{SS} \ll 10^{-7}$ in order for $\tilde{\Lambda}_{DD}$ to be sufficiently large that Higgs portal interactions producing dark neutrinos can be neglected (as we did). This is setup is quite contrived and not really justified by any plausible argument.

There is a much simpler model where the values of the needed values of the effective scales emerge quite naturally.

10.2 Heavy fermion F as mediator

Let us extend the seesaw Lagrangian introducing an heavy fermion doublet F with Yukawa couplings y_I to RH neutrinos,

$$\mathcal{L}_F = \bar{F} (i \not{\partial} - M_F) F - y_I (\bar{F} \phi N_I + \bar{N}_I \phi^\dagger F). \quad (34)$$

At scales much below M_F one can integrate out F obtaining the effective Lagrangian

$$-\mathcal{L}_F^{\text{eff}} = \frac{y_I y_J}{M_F} \bar{N}_I N_J \phi^\dagger \phi, \quad (35)$$




Figure 10: Feynman diagrams with a heavy fermion F as mediator and $I, J = D, S$. Integrating out F , the diagram in the left panel leads to the Feynman diagrams in panel (b) of Fig. 1.

where the RH side coincides with the Anisimov operators with the simple identification $\Lambda = M_F$ and $\lambda'_{IJ} = y_I y_J$. The three Anisimov operators in Eq. (15), Higgs-induced neutrino mixing, source neutrino Higgs portal interactions and dark neutrino Higgs portal interactions, can then be regarded as the low energy effective operators generated by the three diagrams in Fig. 10, respectively.

If we take $M_F \sim M_{\text{GUT}}$, $y_S \sim 1$ and $y_D \sim 10^{-7}$, one can this time immediately reproduce the values $\tilde{\Lambda}_{SS} \sim 10^{16}$ GeV, $\tilde{\Lambda}_{DS} \sim 10^{23}$ GeV and $\tilde{\Lambda}_{DD} \sim 10^{30}$ GeV: the correct values to reproduce the observed DM abundance from Higgs-induced RH neutrino mixing, with source neutrino Higgs portal interactions able to thermalise the source neutrino abundance prior to the onset of the oscillations and with a suppressed contribution to dark neutrino production that we have indeed neglected. Notice that since the three couplings λ_{IJ} are the product of just two Yukawa couplings, it is non trivial that the third comes out automatically satisfying correctly the condition for Higgs portal interactions producing dark neutrinos to be negligible. Moreover, they can be well understood imposing a \mathbb{Z}_2 symmetry under which all particles are even, except the dark neutrino that is odd. In this way the small Yukawa coupling $y_D \sim 10^{-7}$ could be regarded as a small symmetry breaking parameter connecting the visible sector to the dark sector.

11. Summary

The DM puzzle might have a solution to much higher scales than usually considered. Neutrino physics is a good place where to look for such a solution. A high scale RH neutrino playing the role of DM particle requires an extension of the SM beyond the minimal type-I seesaw Lagrangian (able to explain neutrino masses and mixing and the matter-antimatter asymmetry of the universe with leptogenesis). Higgs induced sterile-sterile neutrino mixing provides not only a way to produce the dark neutrinos with the right abundance but it also makes them detectable at neutrino telescopes. Higgs portal interactions for the source neutrino enhance the dark neutrino production and allow to lift the scale of leptogenesis up to 100 TeV. Intriguingly, the IceCube collaboration find an excess

in the neutrino flux at ~ 100 TeV that does not seem easy to explain with astrophysical sources but that can be well explained by RHINO DM decays. Further support comes from multimessenger consideration while a crucial test might come soon by an analysis of anisotropies in the very high energy neutrino flux at IceCube.

Acknowledgments

I acknowledge financial support from the STFC Consolidated Grant ST/T000775/1. I wish to thank Kareem Farrag, Patrick Ludl, Adam Murphy, Sergio Palomares-Ruiz, Rome Samanta and Ye-Ling Zhou for a fruitful collaboration on the RHINO DM model.

References

- [1] For a recent review and references see P. Di Bari, *On the origin of matter in the Universe*, Prog. Part. Nucl. Phys. **122** (2022), 103913 [arXiv:2107.13750 [hep-ph]].
- [2] Y. Akrami *et al.* [Planck Collaboration], *Planck 2018 results. I. Overview and the cosmological legacy of Planck*, arXiv:1807.06205 [astro-ph.CO].
- [3] P. Colin, V. Avila-Reese and O. Valenzuela, *Astrophys. J.* **542** (2000), 622-630 [arXiv:astro-ph/0004115 [astro-ph]].
- [4] P. Minkowski, $\mu \rightarrow e\gamma$ *At A Rate Of One Out Of 1-Billion Muon Decays?*, Phys. Lett. B **67** (1977) 421; T. Yanagida, *Horizontal gauge symmetry and masses of neutrinos*, Conf. Proc. C **7902131** (1979) 95. Proceedings of the Workshop on Unified Theory and Baryon Number of the Universe, eds. O. Sawada and A. Sugamoto (KEK, 1979) p.95; P. Ramond, Invited talk given at Conference: C79-02-25 (Feb 1979) p.265-280, CALT-68-709, *The Family Group in Grand Unified Theories*, hep-ph/9809459; M. Gell-Mann, P. Ramond and R. Slansky, in *Supergravity*, eds. P. van Nieuwenhuizen and D. Freedman (North Holland, Amsterdam, 1979) Conf.Proc. C790927 p.315, PRINT-80-0576; R. Barbieri, D. V. Nanopoulos, G. Morchio and F. Strocchi, *Neutrino Masses in Grand Unified Theories*, Phys. Lett. **90B** (1980) 91. R. N. Mohapatra and G. Senjanovic, *Neutrino Mass and Spontaneous Parity Nonconservation*, Phys. Rev. Lett. **44** (1980) 912.
- [5] M. D’Onofrio, K. Rummukainen and A. Tranberg, *Sphaleron Rate in the Minimal Standard Model*, Phys. Rev. Lett. **113** (2014) no.14, 141602 [arXiv:1404.3565].
- [6] M. Fukugita and T. Yanagida, *Baryogenesis Without Grand Unification*, Phys. Lett. B **174** (1986) 45.
- [7] For a review on thermal leptogenesis see S. Blanchet and P. Di Bari, *The minimal scenario of leptogenesis*, New J. Phys. **14** (2012), 125012 [arXiv:1211.0512 [hep-ph]]. Copy to ClipboardDownload
- [8] S. Dodelson and L. M. Widrow, *Sterile-neutrinos as dark matter*, Phys. Rev. Lett. **72** (1994), 17-20 [arXiv:hep-ph/9303287 [hep-ph]].

- [9] T. Asaka, S. Blanchet and M. Shaposhnikov, *The nuMSM, dark matter and neutrino masses*, Phys. Lett. B **631** (2005), 151-156 [arXiv:hep-ph/0503065 [hep-ph]].
- [10] X. D. Shi and G. M. Fuller, *A New dark matter candidate: Nonthermal sterile neutrinos*, Phys. Rev. Lett. **82** (1999), 2832-2835 [arXiv:astro-ph/9810076 [astro-ph]].
- [11] A. D. Dolgov and S. H. Hansen, *Astropart. Phys.* **16** (2002), 339-344 doi:10.1016/S0927-6505(01)00115-3 [arXiv:hep-ph/0009083 [hep-ph]].
- [12] E. Bulbul, M. Markevitch, A. Foster, R. K. Smith, M. Loewenstein and S. W. Randall, *Detection of An Unidentified Emission Line in the Stacked X-ray spectrum of Galaxy Clusters*, Astrophys. J. **789** (2014), 13 [arXiv:1402.2301 [astro-ph.CO]].
- [13] A. Boyarsky, O. Ruchayskiy, D. Iakubovskiy and J. Franse, *Unidentified Line in X-Ray Spectra of the Andromeda Galaxy and Perseus Galaxy Cluster*, Phys. Rev. Lett. **113** (2014), 251301 [arXiv:1402.4119 [astro-ph.CO]].
- [14] K. N. Abazajian, *Resonantly Produced 7 keV Sterile Neutrino Dark Matter Models and the Properties of Milky Way Satellites*, Phys. Rev. Lett. **112** (2014) no.16, 161303 [arXiv:1403.0954 [astro-ph.CO]].
- [15] K. N. Abazajian, *Technical Comment on "The dark matter interpretation of the 3.5-keV line is inconsistent with blank-sky observations"*, [arXiv:2004.06170 [astro-ph.HE]]; A. Boyarsky, D. Malyshev, O. Ruchayskiy and D. Savchenko, *Technical comment on the paper of Dessert et al. "The dark matter interpretation of the 3.5 keV line is inconsistent with blank-sky observations"*, [arXiv:2004.06601 [astro-ph.CO]]; C. Dessert, N. L. Rodd and B. R. Safdi, *Response to a comment on Dessert et al. "The dark matter interpretation of the 3.5 keV line is inconsistent with blank-sky observations"*, Phys. Dark Univ. **30** (2020), 100656 [arXiv:2006.03974 [astro-ph.CO]]; D. Sicilian, N. Cappelluti, E. Bulbul, F. Civano, M. Moschetti and C. S. Reynolds, *Probing the Milky Way's Dark Matter Halo for the 3.5 keV Line*, [arXiv:2008.02283 [astro-ph.HE]].
- [16] [XRISM Science Team], *Science with the X-ray Imaging and Spectroscopy Mission (XRISM)*, [arXiv:2003.04962 [astro-ph.HE]].
- [17] E. K. Akhmedov, V. A. Rubakov and A. Y. Smirnov, *Baryogenesis via neutrino oscillations*, Phys. Rev. Lett. **81** (1998), 1359-1362 [arXiv:hep-ph/9803255 [hep-ph]].
- [18] T. Asaka and M. Shaposhnikov, *The ν MSM, dark matter and baryon asymmetry of the universe*, Phys. Lett. B **620** (2005), 17-26 [arXiv:hep-ph/0505013 [hep-ph]].
- [19] J. Ghiglieri and M. Laine, *Sterile neutrino dark matter via coinciding resonances*, JCAP **07** (2020), 012 [arXiv:2004.10766 [hep-ph]].
- [20] A. Ariga *et al.* [FASER], *FASER: ForwArd Search ExpeRiment at the LHC*, [arXiv:1901.04468 [hep-ex]].

- [21] A. Anisimov and P. Di Bari, *Cold Dark Matter from heavy Right-Handed neutrino mixing*, Phys. Rev. D **80** (2009), 073017 [arXiv:0812.5085 [hep-ph]].
- [22] P. Di Bari, P. O. Ludl and S. Palomares-Ruiz, *Unifying leptogenesis, dark matter and high-energy neutrinos with right-handed neutrino mixing via Higgs portal*, JCAP **11** (2016), 044 [arXiv:1606.06238 [hep-ph]].
- [23] A. Datta, R. Roshan and A. Sil, *Imprint of the Seesaw Mechanism on Feebly Interacting Dark Matter and the Baryon Asymmetry*, Phys. Rev. Lett. **127** (2021) no.23, 231801 [arXiv:2104.02030 [hep-ph]].
- [24] A. Anisimov, *Majorana Dark Matter*, [arXiv:hep-ph/0612024 [hep-ph]].
- [25] See talk by C. DeGrande in this workshop.
- [26] See talk by Jim Talbert in this workshop.
- [27] B. Patt and F. Wilczek, *Higgs-field portal into hidden sectors*, [arXiv:hep-ph/0605188 [hep-ph]].
- [28] P. Di Bari and A. Murphy, *Completing RHINO*, JHEP **03** (2023), 063 [arXiv:2210.10801 [hep-ph]].
- [29] P. Di Bari, K. Farrag, R. Samanta and Y. L. Zhou, *Density matrix calculation of the dark matter abundance in the Higgs induced right-handed neutrino mixing model*, JCAP **10** (2020), 029 [arXiv:1908.00521 [hep-ph]].
- [30] F. Halzen and D. Hooper, Rept. Prog. Phys. **65** (2002), 1025-1078 doi:10.1088/0034-4885/65/7/201 [arXiv:astro-ph/0204527 [astro-ph]].
- [31] M. G. Aartsen *et al.* [IceCube Collaboration], *First observation of PeV-energy neutrinos with IceCube*, Phys. Rev. Lett. **111** (2013) 021103 [arXiv:1304.5356 [astro-ph.HE]]. M. G. Aartsen *et al.* [IceCube Collaboration], *Science* **342** (2013) 1242856 [arXiv:1311.5238 [astro-ph.HE]]; M. G. Aartsen *et al.* [IceCube Collaboration], Phys. Rev. Lett. **113** (2014) 101101 [arXiv:1405.5303 [astro-ph.HE]].
- [32] C. Haack *et al.* [IceCube], *A measurement of the diffuse astrophysical muon neutrino flux using eight years of IceCube data*, PoS **ICRC2017** (2018), 1005 doi:10.22323/1.301.1005
- [33] M. G. Aartsen *et al.* [IceCube], *The IceCube Neutrino Observatory - Contributions to ICRC 2017 Part II: Properties of the Atmospheric and Astrophysical Neutrino Flux*, [arXiv:1710.01191 [astro-ph.HE]].
- [34] R. Abbasi *et al.* [IceCube], [arXiv:2205.12950 [hep-ex]].
- [35] E. W. Kolb and A. J. Long, *Superheavy dark matter through Higgs portal operators*, Phys. Rev. D **96** (2017) no.10, 103540 [arXiv:1708.04293 [astro-ph.CO]].
- [36] A. Pilaftsis and T. E. J. Underwood, *Resonant leptogenesis*, Nucl. Phys. B **692** (2004), 303-345 [arXiv:hep-ph/0309342 [hep-ph]].