

Status of the operations of CALET for 7.5 years on the International Space Station

T. Tamura,^{*a*,*} Y. Akaike^{*b,c*} and K. Kobayashi^{*b,c*} for the CALET collaboration

^a Kanagawa University, Faculty of Engineering, Department of Applied Physics, 3-27-1 Rokkakubashi, Kanagawa-ku, Yokohama, Japan

^b Waseda University, Waseda Research Institute for Science and Engineering, 17 Kikuicho, Shinjuku, Tokyo, Japan

^c Japan Aerospace Exploration Agency, JEM Utilization Center, Human Spaceflight Technology Directorate,

2-1-1 Sengen, Tsukuba, Ibaraki, Japan

E-mail: ttamura@kanagawa-u.ac.jp

The CALorimetric Electron Telescope (CALET) has successfully been carrying out cosmic-ray observations on the International Space Station since October, 2015. CALET directly measures the cosmic-ray electron spectrum in the energy range of 1 GeV to 20 TeV with a 2 % energy resolution above 30 GeV. In addition, the instrument can measure the spectrum of gamma rays well into the TeV range, and the spectra of protons and nuclei up to a PeV. The scientific operations are implemented by taking into account orbital variations of geomagnetic rigidity cutoff. Scheduled command sequences are used to control the CALET observation modes on orbit. The high-energy (>10 GeV) trigger mode is always active for maintaining maximum exposure to high-energy electrons and other high-energy shower events. Also, around the ISS orbit, calibration data acquisition by, for example, recording pedestal and penetrating particle events, a low-energy electron trigger mode operating at high geomagnetic latitude, a low-energy gamma-ray trigger mode operating at low geomagnetic latitude, and an ultra-heavy trigger mode, are scheduled. As of June 30, 2023, the total observation time is 2818 days with a live time fraction of the total time around 86 %. Nearly 1.86 billion events are collected with the high-energy trigger.

38th International Cosmic Ray Conference (ICRC2023) 26 July - 3 August, 2023 Nagoya, Japan



© Copyright owned by the author(s) under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License (CC BY-NC-ND 4.0).

1. Cosmic-ray observations with CALET on the ISS

The CALorimetric Electron Telescope (CALET), which was launched on August 15, 2015, has been carrying out observations of electrons, protons, nuclei, ultra heavy particles, and gamma-rays on the International Space Station (ISS) since October 13, 2015. As shown in Fig. 1, CALET consists of a main detector CAL (Calorimeter), a gamma ray burst monitor CGBM (CALET Gamma-ray Burst Monitor), and other supporting sensors such as a GPSR (GPS Receiver) and an ASC (Advanced Stellar Camera). CAL consists of a combination of three types of detectors. A CHD (CHarge Detector) placed on the top measures the charge of the incident particles by 28 plastic scintillator bars. An IMC (IMaging Calorimeter) placed in the middle detects directions of incident particles and starting points of showers by 7168 scintillating fibers. A TASC (Total AbSorption Calorimeter) placed at the bottom discriminates kinds of incident particles and determines their energies by 192 PWO (PbWO₄) scintillators. CALET is controlled by an MDC (Mission Data Controller) to collect data from detectors, communicate data with the ground, and execute commands. For about 8 years after launch, CALET has continued its observations smoothly without any major failures. Only one of the 7168 channels of the IMC detector has a permanent failure and only one has a gain drop after launch. Through CALET observations so far, we have obtained a number of results, including an energy spectrum of electrons up to 4.8 TeV [1, 2], energy spectra of nuclei from protons to nickel [3-10], low energy electron precipitation phenomena[11-13], a charge dependence of the solar modulation of cosmic rays[14], gamma-ray burst observation[15], and search of gamma-ray counter part of gravitational wave events[16–18].

2. Operations for CALET

2.1 Normal and Special Operations

Normal operations for CALET include creating schedule files and checking status of the CALET equipment and qualities of its data. The schedule file is created for about 4 days, which is the upper limit of the file size, and updated every Monday, Wednesday, and Friday around 3:00 UT. According




Figure 1: An overview of the CALET instrument




Figure 2: Expected LE-electron count rate

to the regular observation schedule mentioned below, we create a schedule file based on the two-line orbital element (TLE) containing the ISS orbital information obtained from Space-Track[19], and send it to TsuKuba Space Center (TKSC) of JAXA. This creation and submission of the schedule file is done automatically. On the other hand, Japanese shift members take turns checking the CALET DQC (Data Quality Check) website distributed by the WCOC (Waseda CALET Operations Center) every day and reporting the check results by email.

We sometimes need special operations related to ISS operations or CALET malfunctions. We have to deal with tasks related to ISS operations such as reboosts for altitude change, PDAM (Predetermined Debris Avoidance Maneuver), power reduction requests, out gas of other equipment. We also have to handle known events such as MDC auto-reboot by SEU (Single Event Upset), no MDC response for ping due to a filled up ARP (Address Resolution Protocol) table, and so on.

2.2 Observation schedule

CAL has five observation modes, high energy trigger (HE), low energy electron trigger (LEelectron), low energy gamma-ray trigger (LE- γ), pedestal trigger, single trigger (p, He). The HE mode which is set for main purpose of CALET to observe high energy electrons over 10 GeV is always active. Its count rate is less than 10 Hz at any longitude. The LE-electron mode is set to observe low energy electron more than 1 GeV in high latitude of low cutoff rigidities. Because a count rate of the low energy electron observation reaches to around 300 Hz as shown in Fig. 2, each observation period is limited to 90 seconds. The LE- γ mode is set to observe gamma-rays over 1 GeV. In order to limit a count rate of several tens events/second, the LE- γ mode is active in low latitude within plus minus 20 degrees as shown in Fig. 2. We take pedestal data every 23 minutes. Because the orbital period of the ISS is 90 minutes, we expect to acquire the data at all unbiased points. We also take proton or helium events as calibration data by setting single trigger mode at a fixed time (11:30 - 14:30 UT) every day. We can calibrate signal change of each channel of CHD, IMC, and TASC for single particles.




Figure 3: Schedule commands and their execution timing on the ISS track

The photomultiplier tubes (PMTs) of CGBM may be damaged in gain at high count rates. Therefore, the PMTs are protected by turning off the high voltages (HVs) for the PMTs in the high count rate region, such as north and south high altitude regions and South Atlantic Anomaly (SAA). The MDC changes the trigger modes and set the HVs on and off to protect the CGBM PMTs according to commands and execution times written in the schedule file as shown in Fig. 3.

2.3 Data processing operation

The raw data is transferred to the ground in real time during the 80% AOS (Acquisition Of Signal) period. They are recorded in JAXA's TKSC database, and sent to also WCOC. Based on the raw data, WCOC displays an overview of the data and the quality of the data in Quick Look (QL) on web pages, enabling real-time monitoring from anywhere at any time. It allows us to ask the TKSC operator to send a safetying command if necessary.

Data that could not be transferred to the ground in real time during LOS (Loss Of Signal) period will be transferred afterward to the TKSC and recorded in the database. In TKSC, AOS and LOS data are merged into one file as Level 0 data for one hour according to the time series. This Level 0 data file is transferred to the WCOC every hour.

In WCOC, Level 1 data is created by converting Level 0 data into physical quantities and used to create QL on the web pages. Furthermore, higher order data files (such as Level 2) are created for detailed scientific data analysis.




Figure 4: Data processing flow

3. Status of CALET operation

Fig. 5 shows the daily observation time and the actual observation time (live time). Observation time became less than 24 hours when some troubles occurred and the countermeasures were taken, however it can be seen that the observation times were quickly recovered. As shown in Fig. 6 a total observation time of CALET from October 13, 2015 to June 30, 2023 is 2818 days. A live time, which is an actual observation time excluding the dead time for the observation time, is 85% or more. An exposure in the HE trigger mode during the observation time has reached $S\Omega T = 248$ m²sr days. Total data obtained by the HE trigger mode reached 1.86×10^9 events on June 30, 2023. It is steadily increasing as shown in Fig. 7. The spectrum of energy loss at TASC for all 4.05 billion events acquired during the observation time is shown in Fig. 8.




Figure 5: Daily observation time and actual observation time (live time)




Figure 6: Accumulated observation time, actual observation time (live time), and dead time




Figure 7: Accumulated event number of the HE trigger

Figure 8: Energy deposit in TASC

4. Summary

CALET has been carrying out cosmic-ray observations on the space station for more than seven years while maintaining the expected instrument performance. Responses related to ISS operations, such as reboosting and power reduction requests, are being properly implemented in collaboration with JAXA. When malfunctions occur, they are identified by checking observation data, and measures are taken quickly to restore normal observation in cooperation with JAXA.

Acknowledgements

We gratefully acknowledge JAXA contributions to the development of CALET and to the operations onboard the International Space Station. The CALET effort in Italy is supported by ASI under Agreement No. 2013- 018-R.0 and its amendments. The CALET effort in the United States is supported by NASA through Grants No. 80NSSC20K0397, No. 80NSSC20K0399, and No. NNH18ZDA001N-APRA18-0004. This work is supported in part by JSPS Grant-in-Aid for Scientific Research (S) Grant No.19H05608 in Japan.

References

- O. Adriani, Y.Asaoka, S.Torii et al., "Energy Spectrum of Cosmic-Ray Electron and Positron from 10 GeV to 3 TeV Observed with the Calorimetric Electron Telescope on the International Space Station", Phys. Rev. Lett. 119, 181101, 2017
- [2] O. Adriani, Y.Asaoka, S.Torii et al., "Extended Measurements of Cosmic-ray Electron and Positron Spectrum from 11 GeV to 4.8 TeV with the Calorimetric Electron Telescope on the International Space Station", Phys. Rev. Lett. 120, 261102, 2018
- [3] O.Adriani, Y.Asaoka, P.S.Morrecchesi, S.Torii et al., "Direct Measurement of the Cosmic-Ray Proton Spectrum from 50 GeV to 10 TeV with the Calorimetric Electron Telescope on the International Space Station", Physical Review Letters 122, 181102, 2019
- [4] O. Adriani, K. Kobayashi, P.S. Marrocchesi, S. Torii et al., "Observation of Spectral Structures in the Flux of Cosmic-Ray Protons from 50 GeV to 60 TeV with the Calorimetric Electron Telescope on the International Space Station", Physical Review Letters 129, 101102, 2022
- [5] O. Adriani, K. Kobayashi, P. Brogi et al., "Direct Measurement of the Cosmic-Ray Helium Spectrum from 40 GeV to 250 TeV with the Calorimetric Electron Telescope on the International Space Station", Physical Review Letters 130, 171002, 2023
- [6] O. Adriani, Y.Akaike, P. Maestro, et al., "The Cosmic-ray Boron Flux Measured from 8.4 GeV/n to 3.8 TeV/n with the Calorimetric Electron Telescope on the International Space Station", Physical Review Letters 129, 251103, 2022
- [7] P. Maestro et al., "Measurement of cosmic-ray carbon and oxygen energy spectra with CALET", Advances in Space Research 64, 2538-2545, 2019
- [8] O.Adriani, Y.Akaike, P. Maestro et al., "Direct Measurement of the Cosmic-Ray Carbon and Oxygen Spectra from 10 GeV/n to2.2 TeV/n with the Calorimetric Electron Telescope on the International Space Station", Physical Review Letters 125, 251102, 2020
- [9] O.Adriani, Y.Akaike, C. Checchia, et al., "Measurement of the Iron Spectrum in Cosmic Rays from 10 GeV/n to 2.0 TeV/n with the Calorimetric Electron Telescope on the International Space Station", Physical Review Letters 126, 241101 (2021)
- [10] O. Adriani, Y. Akaike, C. Checchia, et al., "Direct Measurement of the Nickel Spectrum in Cosmic Rays in the Energy Range from 8.8 GeV/n to 240 GeV/n with CALET on the International Space Station", Physical Review Letters 128,131103, 2022
- [11] R.Kataoka, Y.Asaoka, S.Torii, T.Terasawa, S.Ozawa, T.Tamura, Y.Shimizu, Y.Akaike, and M.Mori, "Relativistic electron precipitation at International Space Station: Space weather monitoring by Calorimetric Electron Telescope", Geophysical Research Letters, 10.1002/2016GL068930, 2016

- [12] R.Kataoka, Y.Asaoka, S.Torii, S.Nakahira, H.Ueno, S.Miyake, Y.Miyoshi, et al., "Plasma Waves Causing Relativistic Electron Precipitation Events at International Space Station: Lessons From Conjunction Observations With Arase Satellite", Journal of Geophysical Research: Space Physics, 125, e2020JA027875, 2020
- [13] Bruno, L. W. Blum, G. A. de Nolfo, R. Kataoka, S. Torii, A. D. Greeley, S. G. Kanekal, W. Ficklin, T. G. Guzik, S. Nakahira, "EMIC-Wave Driven Electron Precipitation Observed by CALET on the International Space Station", Geophysical Research Letters, Vol.49, Issue 6, e2021GL097529, 2022
- [14] O. Adriani, S. Miyake, K. Munakata, Y. Akaike et al., "Charge-sign dependent cosmic-ray modulation observed with the Calorimetric Electron Telescope on the International Space Station", Physical Review Letters 130, 211001, 2023
- [15] Y.Kawakubo et al. "Detection of the thermal component in GRB 160107A", PASJ 70, 6, 2018
- [16] O. Adriani, Y.Asaoka, S. Nakahira, T.Sakamoto et al., "CALET UPPER LIMITS ON X-RAY AND GAMMA-RAY COUNTERPARTS OF GW 151226", The Astrophysical Journal Letters, 829, L20, 2016
- [17] O. Adriani, Y. Asaoka, M. Mori el al., "Search for GeV Gamma-Ray Counterparts of Gravitational Wave Events by CALET", The Astrophysical Journal, 863, 160, 2018
- [18] O.Adriani, M.L.Cherry, Y.Kawakubo et al., "CALET Search for Electromagnetic Counterparts of Gravitational Waves during the LIGO/Virgo O3 Run", The Astrophysical Journal, 933:85(16pp), 2022
- [19] https://www.space-track.org

Full Author List: CALET Collaboration

O. Adriani^{1,2}, Y. Akaike^{3,4}, K. Asano⁵, Y. Asaoka⁵, E. Berti^{2,6}, G. Bigongiari^{7,8}, W.R. Binns⁹, M. Bongi^{1,2}, P. Brogi^{7,8}, A. Bruno¹⁰, N. Cannady^{11,12,13}, G. Castellini⁶, C. Checchia^{7,8}, M.L. Cherry¹⁴, G. Collazuol^{15,16}, G.A. de Nolfo¹⁰, K. Ebisawa¹⁷, A.W. Ficklin¹⁴, H. Fuke¹⁷, S. Gonzi^{1,2,6}, T.G. Guzik¹⁴, T. Hams¹¹, K. Hibino¹⁸, M. Ichimura¹⁹, K. Ioka²⁰, W. Ishizaki⁵, M.H. Israel⁹, K. Kasahara²¹, J. Kataoka²², R. Kataoka²³, Y. Katayose²⁴, C. Kato²⁵, N. Kawanaka²⁰, Y. Kawakubo¹⁴, K. Kobayashi^{3,4}, K. Kohri²⁶, H.S. Krawczynski⁹, J.F. Krizmanic¹², P. Maestro^{7,8}, P.S. Marrocchesi^{7,8}, A.M. Messineo^{8,27}, J.W. Mitchell¹², S. Miyake²⁸, A.A. Moiseev^{29,12,13}, M. Mori³⁰, N. Mori², H.M. Motz¹⁸, K. Munakata²⁵, S. Nakahira¹⁷, J. Nishimura¹⁷, S. Okuno¹⁸, J.F. Ormes³¹, S. Ozawa³², L. Pacini^{2,6}, P. Papini², B.F. Rauch⁹, S.B. Ricciarini^{2,6}, K. Sakai^{11,12,13}, T. Sakamoto³³, M. Sasaki^{29,12,13}, Y. Shimizu¹⁸, A. Shiomi³⁴, P. Spillantini¹, F. Stolzi^{7,8}, S. Sugita³³, A. Sulaj^{7,8}, M. Takita⁵, T. Tamura¹⁸, T. Terasawa⁵, S. Torii³, Y. Tsunesada^{35,36}, Y. Uchihori³⁷, E. Vannuccini², J.P. Wefel¹⁴, K. Yamaoka³⁸, S. Yanagita³⁹, A. Yoshida³³, K. Yoshida²¹, and W.V. Zober⁹

¹Department of Physics, University of Florence, Via Sansone, 1 - 50019, Sesto Fiorentino, Italy, ²INFN Sezione di Firenze, Via Sansone, 1 - 50019, Sesto Fiorentino, Italy, ³Waseda Research Institute for Science and Engineering, Waseda University, 17 Kikuicho, Shinjuku, Tokyo 162-0044, Japan, ⁴JEM Utilization Center, Human Spaceflight Technology Directorate, Japan Aerospace Exploration Agency, 2-1-1 Sengen, Tsukuba, Ibaraki 305-8505, Japan, ⁵Institute for Cosmic Ray Research, The University of Tokyo, 5-1-5 Kashiwa-no-Ha, Kashiwa, Chiba 277-8582, Japan, ⁶Institute of Applied Physics (IFAC), National Research Council (CNR), Via Madonna del Piano, 10, 50019, Sesto Fiorentino, Italy, ⁷Department of Physical Sciences, Earth and Environment, University of Siena, via Roma 56, 53100 Siena, Italy, ⁸INFN Sezione di Pisa, Polo Fibonacci, Largo B. Pontecorvo, 3 - 56127 Pisa, Italy, ⁹Department of Physics and McDonnell Center for the Space Sciences, Washington University, One Brookings Drive, St. Louis, Missouri 63130-4899, USA, ¹⁰Heliospheric Physics Laboratory, NASA/GSFC, Greenbelt, Maryland 20771, USA, ¹¹Center for Space Sciences and Technology, University of Maryland, Baltimore County, 1000 Hilltop Circle, Baltimore, Maryland 21250, USA, ¹²Astroparticle Physics Laboratory, NASA/GSFC, Greenbelt, Maryland 20771, USA, ¹³Center for Research and Exploration in Space Sciences and Technology, NASA/GSFC, Greenbelt, Maryland 20771, USA, ¹⁴Department of Physics and Astronomy, Louisiana State University, 202 Nicholson Hall, Baton Rouge, Louisiana 70803, USA, ¹⁵Department of Physics and Astronomy, University of Padova, Via Marzolo, 8, 35131 Padova, Italy, ¹⁶INFN Sezione di Padova, Via Marzolo, 8, 35131 Padova, Italy, ¹⁷Institute of Space and Astronautical Science, Japan Aerospace Exploration Agency, 3-1-1 Yoshinodai, Chuo, Sagamihara, Kanagawa 252-5210, Japan, ¹⁸Kanagawa University, 3-27-1 Rokkakubashi, Kanagawa, Yokohama, Kanagawa 221-8686, Japan, ¹⁹Faculty of Science and Technology, Graduate School of Science and Technology, Hirosaki University, 3, Bunkyo, Hirosaki, Aomori 036-8561, Japan, ²⁰Yukawa Institute for Theoretical Physics, Kyoto University, Kitashirakawa Oiwake-cho, Sakyo-ku, Kyoto, 606-8502, Japan, ²¹Department of Electronic Information Systems, Shibaura Institute of Technology, 307 Fukasaku, Minuma, Saitama 337-8570, Japan, ²²School of Advanced Science and Engineering, Waseda University, 3-4-1 Okubo, Shinjuku, Tokyo 169-8555, Japan, ²³National Institute of Polar Research, 10-3, Midori-cho, Tachikawa, Tokyo 190-8518, Japan, ²⁴Faculty of Engineering, Division of Intelligent Systems Engineering, Yokohama National University, 79-5 Tokiwadai, Hodogaya, Yokohama 240-8501, Japan, ²⁵Faculty of Science, Shinshu University, 3-1-1 Asahi, Matsumoto, Nagano 390-8621, Japan, ²⁶Institute of Particle and Nuclear Studies, High Energy Accelerator Research Organization, 1-1 Oho, Tsukuba, Ibaraki, 305-0801, Japan, 27 University of Pisa, Polo Fibonacci, Largo B. Pontecorvo, 3 - 56127 Pisa, Italy, 28 Department of Electrical and Electronic Systems Engineering, National Institute of Technology (KOSEN), Ibaraki College, 866 Nakane, Hitachinaka, Ibaraki 312-8508, Japan, ²⁹Department of Astronomy, University of Maryland, College Park, Maryland 20742, USA, ³⁰Department of Physical Sciences, College of Science and Engineering, Ritsumeikan University, Shiga 525-8577, Japan, ³¹Department of Physics and Astronomy, University of Denver, Physics Building, Room 211, 2112 East Wesley Avenue, Denver, Colorado 80208-6900, USA, 32 Quantum ICT Advanced Development Center, National Institute of Information and Communications Technology, 4-2-1 Nukui-Kitamachi, Koganei, Tokyo 184-8795, Japan, ³³College of Science and Engineering, Department of Physics and Mathematics, Aoyama Gakuin University, 5-10-1 Fuchinobe, Chuo, Sagamihara, Kanagawa 252-5258, Japan, ³⁴College of Industrial Technology, Nihon University, 1-2-1 Izumi, Narashino, Chiba 275-8575, Japan, ³⁵Graduate School of Science, Osaka Metropolitan University, Sugimoto, Sumiyoshi, Osaka 558-8585, Japan, ³⁶Nambu Yoichiro Institute for Theoretical and Experimental Physics, Osaka Metropolitan University, Sugimoto, Sumiyoshi, Osaka 558-8585, Japan, ³⁷National Institutes for Quantum and Radiation Science and Technology, 4-9-1 Anagawa, Inage, Chiba 263-8555, Japan, ³⁸Nagoya University, Furo, Chikusa, Nagoya 464-8601, Japan, 39 College of Science, Ibaraki University, 2-1-1 Bunkyo, Mito, Ibaraki 310-8512, Japan