

## Status and Perspectives on Axion Searches

---

**Maurizio Giannotti**<sup>a,b,\*</sup>

<sup>a</sup>*Centro de Astropartículas y Física de Altas Energías, University of Zaragoza, Zaragoza, 50009, Aragón, Spain*

<sup>b</sup>*Physical Sciences, Barry University, 11300 NE 2nd Ave., Miami Shores, FL 33161, USA*

*E-mail:* [mgiannotti@unizar.es](mailto:mgiannotti@unizar.es)

The search for axions and axion-like particles (ALPs) remains a major endeavor in modern physics investigation. Axions play essential roles in the quest to understand dark matter, the strong CP problem, and various astrophysical phenomena. This paper provides a very brief overview of the current status of experimental efforts, highlighting significant advancements, ongoing projects, and future opportunities. Particular attention is given to cavity haloscopes, helioscopes, and laboratory-based light-shining-through-wall experiments, as well as astrophysical probes. Some future perspectives are also discussed.

*2nd General Meeting of COST Action COSMIC WISPer  
3-6 September 2024  
Istanbul, Turkey*

---

\*Speaker

## 1. Introduction

Axions and, more generally, Axion-Like Particles (ALPs) are prominent in modern theoretical physics as they emerge in many extensions of the Standard Model (SM) [1]. They are generally defined as pseudoscalar particles described by the low-energy Lagrangian:

$$\mathcal{L}_a = \frac{1}{2}(\partial_\mu a)^2 - m_a^2 a^2 - \sum_{f=e,p,n} g_{af} a \bar{\psi}_f \gamma_5 \psi_f - \frac{1}{4} g_{a\gamma} a F_{\mu\nu} \tilde{F}^{\mu\nu}, \quad (1)$$

where  $a$  is the ALP field,  $m_a$  its mass,  $g_{af}$  and  $g_{a\gamma}$  parameterize its couplings to fermions ( $\psi_f$ ) and photons, and  $F$  and  $\tilde{F}$  represent the electromagnetic field tensor and its dual. The most prominent example of ALP is the QCD axion, a key prediction of the Peccei-Quinn solution to the strong CP problem [2–4] and a favored dark matter (DM) candidate [5–7]. Beyond QCD axions, ALPs naturally emerge in other SM extensions, such as string theory [8–10], and often exhibit weaker constraints, lacking specific mass-coupling relations.

The experimental search for axions and ALPs has seen a significant resurgence, driven by innovative techniques capable of probing previously unexplored regions of the parameter space [11–13]. Many of these regions are motivated by astrophysical and cosmological considerations, making the prospects for near-term discovery particularly promising [14–16]. This review provides a concise overview of the recent progress in axion and ALP research, with a focus on the axion-photon coupling, while highlighting key developments and near-future opportunities. For technical details, readers may refer to the comprehensive discussion in Ref. [11], and for a broader set of references, to the informative webpage [17].

## 2. Current Experimental Landscape

Axion searches through the axion-photon coupling are broadly divided into 3 categories, depending on the specific axion source:

- *Pure laboratory experiments*, such as light shining through a wall (LSW). In this case, axions are produced in a laboratory setting. These searches are appealing due to the complete control over the source and minimal assumptions about production mechanisms. However, their sensitivity is limited due to the highly suppressed signal ( $\propto g_{a\gamma}^4$ ).
- *Helioscopes*. In this case, the source (the Sun) is well understood, requiring minimal assumptions about axion production [18]. An exception is axion production in the solar magnetic field, which relies on much less well-established assumptions [19–21].
- *Haloscopes*. They search for dark matter axions. Thus, the sensitivity to the axion coupling is necessarily dependent on the specific assumption about the fraction of dark matter made up by axions, a fact that is not known and that cannot be deduced by current theoretical considerations [7].

### 2.1 Laboratory Searches

The main protagonists are the LSW experiments, in which axions are produced from a coherent source of photons (Light) interacting with a magnetic field, and detected as photons (Shining) after

**Table 1:** Most relevant LSW experiments, with some technical considerations. Data from Ref. [11].

Experiment	$B(\text{T})$	$L(\text{m})$	$g_{a\gamma} [\text{GeV}^{-1}]$	Notes
ALPS-I [22]	5	4.3	$5 \times 10^{-8}$	completed
CROWS [23]	3	0.15	$9.9 \times 10^{-8}$	uses microwaves
OSQAR [24]	9	14.3	$3.5 \times 10^{-8}$	dipole magnet from LHC
ALPS-II [25]	5	100	$2 \times 10^{-11}$	dipole magnet from HERA
JURA [28]	13	426	$10^{-12}$	concept

being reconverted after passing through a region opaque to photons (Wall). The forefront LSW experiment is ALPS II [26], in DESY, which is expected to probe the axion-photon coupling down to  $g_{a\gamma} \approx 2 \times 10^2 \text{ GeV}^{-1}$ , for masses below 0.1 meV or so (see Tab. 1). The first run finished on May 2024 and a related publication is expected for early 2025 [27]. The full optics is expected to be completed in 2025 [27]. Ongoing discussions within the Physics Beyond Colliders (PBC) working group at CERN are considering the possibility of a larger scale LSW experiment, JURA, which would significantly increase the magnetic field length while bringing its intensity up to 13 T [28].

Polarization experiments operate on a concept similar to LSW, but focusing on detecting anomalies in light polarization caused by axions or other WISPs. The prototype experiment in this category is PVLAS [29, 30], while an improved version, VMB@CERN, is currently under discussion within the PBC [31]. However, the feasibility of this upgraded experiment remains uncertain.

## 2.2 Solar axion detection

Currently, the optimal way to detect solar axions is the so called Sikivie Helioscope [32], which relies on the conversion of solar axions into photons in a terrestrial magnetic field. An advantage of this method with respect to pure laboratory experiments, is the possibility to measure the axion flux produced through different mechanisms, even those not induced by the axion-photon coupling (see, e.g., [33]), as well as the flux of other WISPs [34, 35].

The currently most advanced example is the CERN Axion Solar Telescope (CAST), which recently published updated analysis results [36], pushing the limits on the axion-photon coupling below any other experimental results, and probing a wide mass region so far accessible only to astrophysics [16, 37, 38]. CAST successor, BabyIAXO [39, 40], a prototype of the future full International Axion Observatory (IAXO) [41], is planned for construction at DESY and expected to begin operations soon, probing significant new regions of the axion parameter space (Cf. left panel of Fig.1).

There are other possibilities to look for solar axions, including using large underground detectors, which, however, have limited chances to compete with the very strong astrophysical bounds [42, 43] in the near future (see, e.g., Ref. [16] for an overview and for references). Another compelling option is to make use of the solar magnetic field rather than a terrestrial one, and to detect the produced photons flux with a space-borne x-ray detector. An explicit analysis, using the NuSTAR detector, was recently presented in Ref. [44]. The advantage of this method relies on the significantly longer magnetic field, which pushes the sensitivity to couplings considerably below the CAST reach, though for a smaller mass range.



### 3. The Road Ahead

Experimental searches for axions and ALPs have made significant strides in recent years, with numerous new experiments proposed and expected to begin within this decade. The current proliferation of experimental proposals brings substantial optimism for exploring a large portion of the axion parameter space in the near future. The outlook is particularly promising for haloscope searches. Projects like MADMAX [57, 58], CADEX [59], and ultra-low-temperature cavity experiments at IBS/CAPP (CULTASK)[60] are set to probe the axion and ALP mass range from the  $\mu\text{eV}$  up to a few 0.1 meV. Tunable axion plasma haloscopes [61], which leverage the axion-plasmon coupling, complement these efforts in the same mass region. Meanwhile, proposals such as TOORAD [62, 63] and BREAD [64] aim to reach even higher masses, extending the search into the meV regime.

Significant progress has also been made in investigating lower mass regions. The KLASH proposal [65], initially designed to target axion masses in the range  $m_a \in [0.3 - 1]\mu\text{eV}$ , has evolved into FLASH [66], which explores slightly higher masses due to changes in the magnet configuration [67]. FLASH is poised to investigate not only the axion parameter space but also other WISPs and gravitational waves [68, 69]. Additionally, ongoing experiments like ABRACADABRA are expected to achieve high sensitivity, potentially reaching the QCD axion band in the sub- $\mu\text{eV}$  range [70]. Laboratory searches are also progressing. ALPS II [71] and upcoming axion helioscopes, including BabyIAXO [39, 40, 72] and IAXO [41], aim to explore regions of parameter space below current astrophysical bounds.

What era could an axion discovery usher in? The discovery of an ALP would not only unveil a new particle and physics at a novel energy scale but also have profound phenomenological implications. Determining an axion mass could facilitate its detection through other experimental approaches, particularly using highly sensitive cavity searches, if the mass falls within an accessible range. If a cavity experiment were to confirm a signal previously identified in a laboratory or solar search, it could precisely determine the ALP dark matter fraction—a groundbreaking result.<sup>1</sup>

Moreover, if the axion couplings turn out to be strong enough to be detectable by IAXO—or potentially even BabyIAXO—the particle’s interactions could have measurable impacts on stellar evolution [74–76]. In this case, axions could also serve as astrophysical messengers, offering unprecedented insights into stellar interiors through methods inaccessible to traditional telescopes [16]. These includes exploring solar properties like temperature [77], metallicity [78] and magnetic field profiles [21, 79], facilitated by anticipated technological advancements.

Beyond the Sun, axions offer remarkable opportunities for studying other stars. They could serve as invaluable tools for examining supergiants, providing insights into their evolutionary stages, core temperatures, and other parameters inaccessible through conventional methods [80, 81]. Similarly, axions might help explore supernova physics, potentially permitting a deeper understanding of the equation of state of nuclear matter [82–88].

<sup>1</sup>Knowing the mass could also enhance LSW experiment sensitivities beyond their nominal capabilities [73].

#### 4. Conclusions

Axion research has made remarkable progress, with significant advancements across experimental, theoretical, and observational fronts. The upcoming decade holds immense potential for breakthroughs, making axion searches one of the most exciting fields in modern physics.

The diversity of experimental approaches—ranging from haloscopes and helioscopes to laboratory based techniques—ensures a comprehensive exploration of the parameter space. These methods are complemented by astrophysical probes that provide critical constraints and opportunities to uncover axion-related phenomena in stellar environments. The synergy between these techniques promises not only to refine our understanding of axion properties but also to push the boundaries of sensitivity to previously unexplored regions.

The next generation of experiments are poised to revolutionize the field. By leveraging cutting-edge technology and innovative experimental designs, these efforts aim to probe deeper into the axion parameter space and potentially reveal the particle’s role in the dark matter puzzle, the strong CP problem, and stellar physics.

Beyond the technical milestones, an axion discovery would mark the dawn of a new era in physics. Such a breakthrough would unveil physics beyond the Standard Model, opening the door to understanding fundamental questions about the universe. Moreover, it would provide a unique tool to investigate astrophysical phenomena, offering insights into stellar interiors, supernova dynamics, and stellar magnetic field structures that are otherwise inaccessible.

The progress achieved so far underscores the transformative potential of axion research. As we move forward, the collaborative efforts of the scientific community, supported by advancements in technology and theoretical understanding, bring us closer to unraveling one of the most profound mysteries in modern science.

#### Acknowledgments

I express my profound gratitude to the local organizing committee for their excellent organization of the 2nd General Meeting of the COST Action CA21106, at the Istinye University, in Istanbul. I acknowledge support from the Spanish Agencia Estatal de Investigación through grant PID2019-108122GB-C31, funded by MCIN/AEI/10.13039/501100011033, and the “European Union NextGenerationEU/PRTR” (Planes complementarios, Programa de Astrofísica y Física de Altas Energías). Additional support comes from grant PGC2022-126078NB-C21, “Aún más allá de los modelos estándar,” funded by MCIN/AEI/10.13039/501100011033 and “ERDF A way of making Europe,” as well as the European Union’s Horizon 2020 research and innovation programme under the European Research Council (ERC) grant agreement ERC-2017-AdG788781 (IAXO+). This work is also based on contributions from COST Action COSMIC WISPerS CA21106, supported by COST (European Cooperation in Science and Technology).

#### References

- [1] A. Ringwald, [arXiv:1407.0546 [hep-ph]].
- [2] S. Weinberg, Phys. Rev. Lett. **40**, 223-226 (1978) doi:10.1103/PhysRevLett.40.223

- [3] F. Wilczek, Phys. Rev. Lett. **40**, 279-282 (1978) doi:10.1103/PhysRevLett.40.279
- [4] R. D. Peccei and H. R. Quinn, Phys. Rev. Lett. **38**, 1440-1443 (1977) doi:10.1103/PhysRevLett.38.1440
- [5] D. J. E. Marsh, Phys. Rept. **643**, 1-79 (2016) doi:10.1016/j.physrep.2016.06.005 [arXiv:1510.07633 [astro-ph.CO]].
- [6] L. Di Luzio, M. Giannotti, E. Nardi and L. Visinelli, Phys. Rept. **870**, 1-117 (2020) doi:10.1016/j.physrep.2020.06.002 [arXiv:2003.01100 [hep-ph]].
- [7] C. B. Adams, N. Aggarwal, A. Agrawal, R. Balafendiev, C. Bartram, M. Baryakhtar, H. Bekker, P. Belov, K. K. Berggren and A. Berlin, *et al.* [arXiv:2203.14923 [hep-ex]].
- [8] P. Svrcek and E. Witten, JHEP **06**, 051 (2006) doi:10.1088/1126-6708/2006/06/051 [arXiv:hep-th/0605206 [hep-th]].
- [9] A. Arvanitaki, S. Dimopoulos, S. Dubovsky, N. Kaloper and J. March-Russell, Phys. Rev. D **81**, 123530 (2010) doi:10.1103/PhysRevD.81.123530 [arXiv:0905.4720 [hep-th]].
- [10] M. Cicoli, M. Goodsell and A. Ringwald, JHEP **10**, 146 (2012) doi:10.1007/JHEP10(2012)146 [arXiv:1206.0819 [hep-th]].
- [11] I. G. Irastorza and J. Redondo, Prog. Part. Nucl. Phys. **102**, 89-159 (2018) doi:10.1016/j.pnpnp.2018.05.003 [arXiv:1801.08127 [hep-ph]].
- [12] P. Sikivie, Rev. Mod. Phys. **93**, no.1, 015004 (2021) doi:10.1103/RevModPhys.93.015004 [arXiv:2003.02206 [hep-ph]].
- [13] I. G. Irastorza, SciPost Phys. Lect. Notes **45**, 1 (2022) doi:10.21468/SciPostPhysLectNotes.45 [arXiv:2109.07376 [hep-ph]].
- [14] M. Giannotti, Nature Phys. **13**, no.6, 530-531 (2017) doi:10.1038/nphys4139
- [15] M. Giannotti, J. Phys. Conf. Ser. **2502**, no.1, 012003 (2023) doi:10.1088/1742-6596/2502/1/012003 [arXiv:2205.06831 [hep-ph]].
- [16] P. Carenza, M. Giannotti, J. Isern, A. Mirizzi and O. Straniero, [arXiv:2411.02492 [hep-ph]].
- [17] C. O'Hare, "cajohare/AxionLimits: AxionLimits," Zenodo, version v1.0, July 2020. DOI: 10.5281/zenodo.3932430. Available at: <https://cajohare.github.io/AxionLimits/>.
- [18] S. Hoof, J. Jaeckel and L. J. Thormaehlen, JCAP **09**, 006 (2021) doi:10.1088/1475-7516/2021/09/006 [arXiv:2101.08789 [hep-ph]].
- [19] E. Guarini, P. Carenza, J. Galan, M. Giannotti and A. Mirizzi, Phys. Rev. D **102**, no.12, 123024 (2020) doi:10.1103/PhysRevD.102.123024 [arXiv:2010.06601 [hep-ph]].
- [20] A. Caputo, A. J. Millar and E. Vitagliano, Phys. Rev. D **101**, no.12, 123004 (2020) doi:10.1103/PhysRevD.101.123004 [arXiv:2005.00078 [hep-ph]].



- [21] C. A. J. O’Hare, A. Caputo, A. J. Millar and E. Vitagliano, *Phys. Rev. D* **102**, no.4, 043019 (2020) doi:10.1103/PhysRevD.102.043019 [arXiv:2006.10415 [astro-ph.CO]].
- [22] K. Ehret, M. Frede, S. Ghazaryan, M. Hildebrandt, E. A. Knabbe, D. Kracht, A. Lindner, J. List, T. Meier and N. Meyer, *et al.* *Phys. Lett. B* **689**, 149-155 (2010) doi:10.1016/j.physletb.2010.04.066 [arXiv:1004.1313 [hep-ex]].
- [23] M. Betz, F. Caspers, M. Gasior, M. Thumm and S. W. Rieger, *Phys. Rev. D* **88**, no.7, 075014 (2013) doi:10.1103/PhysRevD.88.075014 [arXiv:1310.8098 [physics.ins-det]].
- [24] R. Ballou *et al.* [OSQAR], *Phys. Rev. D* **92**, no.9, 092002 (2015) doi:10.1103/PhysRevD.92.092002 [arXiv:1506.08082 [hep-ex]].
- [25] M. D. Ortiz, J. Gleason, H. Grote, A. Hallal, M. T. Hartman, H. Hollis, K. S. Isleif, A. James, K. Karan and T. Kozlowski, *et al.* *Phys. Dark Univ.* **35**, 100968 (2022) doi:10.1016/j.dark.2022.100968 [arXiv:2009.14294 [physics.optics]].
- [26] A. Spector [ALPS], doi:10.3204/DESY-PROC-2009-03/Spector\_Aaron [arXiv:1611.05863 [physics.ins-det]].
- [27] A. Lindner, “Any Light Particle Search II: status and first results,” Colloquium at Heidelberg University, 9 July 2024. Available at: <https://www.physik.uni-heidelberg.de/hephysto/index.php?s=talk&id=11022>.
- [28] A. Lindner, talk given at Physics Beyond Colliders Annual Workshop, CERN, November 2017. Available at: <https://indico.cern.ch/event/644287/>.
- [29] F. Della Valle, A. Ejlli, U. Gastaldi, G. Messineo, E. Milotti, R. Pengo, G. Ruoso and G. Zavattini, *Eur. Phys. J. C* **76**, no.1, 24 (2016) doi:10.1140/epjc/s10052-015-3869-8 [arXiv:1510.08052 [physics.optics]].
- [30] A. Ejlli, F. Della Valle, U. Gastaldi, G. Messineo, R. Pengo, G. Ruoso and G. Zavattini, *Phys. Rept.* **871**, 1-74 (2020) doi:10.1016/j.physrep.2020.06.001 [arXiv:2005.12913 [physics.optics]].
- [31] S. Kunc [VMB@CERN], PoS **COSMICWISPers**, 032 (2024) doi:10.22323/1.454.0032
- [32] P. Sikivie, *Phys. Rev. Lett.* **51**, 1415-1417 (1983) [erratum: *Phys. Rev. Lett.* **52**, 695 (1984)] doi:10.1103/PhysRevLett.51.1415
- [33] L. Di Luzio, J. Galan, M. Giannotti, I. G. Irastorza, J. Jaeckel, A. Lindner, J. Ruz, U. Schneekloth, L. Sohl and L. J. Thormaehlen, *et al.* *Eur. Phys. J. C* **82**, no.2, 120 (2022) doi:10.1140/epjc/s10052-022-10061-1 [arXiv:2111.06407 [hep-ph]].
- [34] T. O’Shea, A. C. Davis, M. Giannotti, S. Vagnozzi, L. Visinelli and J. K. Vogel, *Phys. Rev. D* **110**, no.6, 063027 (2024) doi:10.1103/PhysRevD.110.063027 [arXiv:2406.01691 [hep-ph]].
- [35] T. O’Shea, M. Giannotti, I. G. Irastorza, L. M. Plasencia, J. Redondo, J. Ruz and J. K. Vogel, *JCAP* **06**, 070 (2024) doi:10.1088/1475-7516/2024/06/070 [arXiv:2312.10150 [hep-ph]].



- [36] K. Altenmüller *et al.* [CAST], Phys. Rev. Lett. **133**, no.22, 221005 (2024) doi:10.1103/PhysRevLett.133.221005 [arXiv:2406.16840 [hep-ex]].
- [37] A. Ayala, I. Domínguez, M. Giannotti, A. Mirizzi and O. Straniero, Phys. Rev. Lett. **113**, no.19, 191302 (2014) doi:10.1103/PhysRevLett.113.191302 [arXiv:1406.6053 [astro-ph.SR]].
- [38] M. J. Dolan, F. J. Hiskens and R. R. Volkas, JCAP **10**, 096 (2022) doi:10.1088/1475-7516/2022/10/096 [arXiv:2207.03102 [hep-ph]].
- [39] A. Abeln *et al.* [IAXO], JHEP **05**, 137 (2021) doi:10.1007/JHEP05(2021)137 [arXiv:2010.12076 [physics.ins-det]].
- [40] S. Ahyoune *et al.* [IAXO], [arXiv:2411.13915 [hep-ex]].
- [41] E. Armengaud *et al.* [IAXO], JCAP **06**, 047 (2019) doi:10.1088/1475-7516/2019/06/047 [arXiv:1904.09155 [hep-ph]].
- [42] O. Straniero, C. Pallanca, E. Dalessandro, I. Dominguez, F. R. Ferraro, M. Giannotti, A. Mirizzi and L. Piersanti, Astron. Astrophys. **644**, A166 (2020) doi:10.1051/0004-6361/202038775 [arXiv:2010.03833 [astro-ph.SR]].
- [43] F. Capozzi and G. Raffelt, Phys. Rev. D **102**, no.8, 083007 (2020) doi:10.1103/PhysRevD.102.083007 [arXiv:2007.03694 [astro-ph.SR]].
- [44] J. Ruz, E. Todarello, J. K. Vogel, M. Giannotti, B. Grefenstette, H. S. Hudson, I. G. Hannah, I. G. Irastorza, C. S. Kim and T. O’Shea, *et al.* [arXiv:2407.03828 [astro-ph.CO]].
- [45] N. Du *et al.* [ADMX], Phys. Rev. Lett. **120**, no.15, 151301 (2018) doi:10.1103/PhysRevLett.120.151301 [arXiv:1804.05750 [hep-ex]].
- [46] T. Braine *et al.* [ADMX], Phys. Rev. Lett. **124**, no.10, 101303 (2020) doi:10.1103/PhysRevLett.124.101303 [arXiv:1910.08638 [hep-ex]].
- [47] C. Bartram *et al.* [ADMX], Phys. Rev. Lett. **127**, no.26, 261803 (2021) doi:10.1103/PhysRevLett.127.261803 [arXiv:2110.06096 [hep-ex]].
- [48] G. Rybka, “The US axion searches program, ADMX plans,” presented at the GGI Workshop “Axions across boundaries between Particle Physics, Astrophysics, Cosmology and forefront Detection Technologies,” Florence, Italy, 6 June 2023. Available at: <https://www.ggi.infn.it/showevent.pl?id=439>.
- [49] A. K. Yi, S. Ahn, Ç. Kutlu, J. Kim, B. R. Ko, B. I. Ivanov, H. Byun, A. F. van Loo, S. Park and J. Jeong, *et al.* Phys. Rev. Lett. **130**, no.7, 071002 (2023) doi:10.1103/PhysRevLett.130.071002 [arXiv:2210.10961 [hep-ex]].
- [50] B. M. Brubaker, L. Zhong, Y. V. Gurevich, S. B. Cahn, S. K. Lamoreaux, M. Simanovskaia, J. R. Root, S. M. Lewis, S. Al Kenany and K. M. Backes, *et al.* Phys. Rev. Lett. **118**, no.6, 061302 (2017) doi:10.1103/PhysRevLett.118.061302 [arXiv:1610.02580 [astro-ph.CO]].

- [51] D. Alesini, C. Braggio, G. Carugno, N. Crescini, D. D’Agostino, D. Di Gioacchino, R. Di Vora, P. Falferi, U. Gambardella and C. Gatti, *et al.* Phys. Rev. D **103**, no.10, 102004 (2021) doi:10.1103/PhysRevD.103.102004 [arXiv:2012.09498 [hep-ex]].
- [52] A. Á. Melcón *et al.* [CAST], JHEP **21**, 075 (2020) doi:10.1007/JHEP10(2021)075 [arXiv:2104.13798 [hep-ex]].
- [53] B. T. McAllister, G. Flower, J. Kruger, E. N. Ivanov, M. Goryachev, J. Bourhill and M. E. Tobar, Phys. Dark Univ. **18**, 67-72 (2017) doi:10.1016/j.dark.2017.09.010 [arXiv:1706.00209 [physics.ins-det]].
- [54] J. A. Devlin, M. J. Borchert, S. Erlewein, M. Fleck, J. A. Harrington, B. Latacz, J. Warncke, E. Wursten, M. A. Bohman and A. H. Mooser, *et al.* Phys. Rev. Lett. **126**, no.4, 041301 (2021) doi:10.1103/PhysRevLett.126.041301 [arXiv:2101.11290 [astro-ph.CO]].
- [55] A. V. Gramolin, D. Aybas, D. Johnson, J. Adam and A. O. Sushkov, Nature Phys. **17**, no.1, 79-84 (2021) doi:10.1038/s41567-020-1006-6 [arXiv:2003.03348 [hep-ex]].
- [56] J. L. Ouellet, C. P. Salemi, J. W. Foster, R. Henning, Z. Bogorad, J. M. Conrad, J. A. Formaggio, Y. Kahn, J. Minervini and A. Radovinsky, *et al.* Phys. Rev. Lett. **122**, no.12, 121802 (2019) doi:10.1103/PhysRevLett.122.121802 [arXiv:1810.12257 [hep-ex]].
- [57] A. Caldwell *et al.* [MADMAX Working Group], Phys. Rev. Lett. **118**, no.9, 091801 (2017) doi:10.1103/PhysRevLett.118.091801 [arXiv:1611.05865 [physics.ins-det]].
- [58] P. Brun *et al.* [MADMAX], Eur. Phys. J. C **79**, no.3, 186 (2019) doi:10.1140/epjc/s10052-019-6683-x [arXiv:1901.07401 [physics.ins-det]].
- [59] B. Aja, S. A. Cuendis, I. Arregui, E. Artal, R. Belén Barreiro, F. J. Casas, M. C. de Ory, A. Díaz-Morcillo, L. de la Fuente and J. D. Gallego, *et al.* JCAP **11**, 044 (2022) doi:10.1088/1475-7516/2022/11/044 [arXiv:2206.02980 [hep-ex]].
- [60] Y. K. Semertzidis, J. E. Kim, S. Youn, J. Choi, W. Chung, S. Haciomeroglu, D. Kim, J. Kim, B. Ko and O. Kwon, *et al.* [arXiv:1910.11591 [physics.ins-det]].
- [61] M. Lawson, A. J. Millar, M. Pancaldi, E. Vitagliano and F. Wilczek, Phys. Rev. Lett. **123**, no.14, 141802 (2019) doi:10.1103/PhysRevLett.123.141802 [arXiv:1904.11872 [hep-ph]].
- [62] D. J. E. Marsh, K. C. Fong, E. W. Lentz, L. Smejkal and M. N. Ali, Phys. Rev. Lett. **123**, no.12, 121601 (2019) doi:10.1103/PhysRevLett.123.121601 [arXiv:1807.08810 [hep-ph]].
- [63] J. Schütte-Engel, D. J. E. Marsh, A. J. Millar, A. Sekine, F. Chadha-Day, S. Hoof, M. N. Ali, K. C. Fong, E. Hardy and L. Šmejkal, JCAP **08**, 066 (2021) doi:10.1088/1475-7516/2021/08/066 [arXiv:2102.05366 [hep-ph]].
- [64] J. Liu *et al.* [BREAD], Phys. Rev. Lett. **128**, no.13, 131801 (2022) doi:10.1103/PhysRevLett.128.131801 [arXiv:2111.12103 [physics.ins-det]].

- [65] D. Alesini, D. Babusci, P. Beltrame, S.J., F. Björkeröth, F. Bossi, P. Ciambone, G. Delle Monache, D. Di Gioacchino, P. Falferi and A. Gallo, *et al.* [arXiv:1911.02427 [physics.ins-det]].
- [66] D. Alesini, D. Babusci, P. Beltrame, F. Bossi, P. Ciambone, A. D’Elia, D. Di Gioacchino, G. Di Pirro, B. Döbrich and P. Falferi, *et al.* Phys. Dark Univ. **42**, 101370 (2023) doi:10.1016/j.dark.2023.101370 [arXiv:2309.00351 [physics.ins-det]].
- [67] C. Gatti, P. Gianotti, C. Ligi, M. Raggi and P. Valente, Universe **7**, no.7, 236 (2021) doi:10.3390/universe7070236
- [68] C. Gatti, L. Visinelli and M. Zantedeschi, Phys. Rev. D **110**, no.2, 023018 (2024) doi:10.1103/PhysRevD.110.023018 [arXiv:2403.18610 [gr-qc]].
- [69] L. Visinelli, [arXiv:2411.19441 [astro-ph.CO]].
- [70] Y. Kahn, B. R. Safdi and J. Thaler, Phys. Rev. Lett. **117**, no.14, 141801 (2016) doi:10.1103/PhysRevLett.117.141801 [arXiv:1602.01086 [hep-ph]].
- [71] R. Bähre, B. Döbrich, J. Dreyling-Eschweiler, S. Ghazaryan, R. Hodajerdi, D. Horns, F. Januschek, E. A. Knabbe, A. Lindner and D. Notz, *et al.* JINST **8**, T09001 (2013) doi:10.1088/1748-0221/8/09/T09001 [arXiv:1302.5647 [physics.ins-det]].
- [72] T. Dafni and J. Galán, Universe **8**, no.1, 37 (2022) doi:10.3390/universe8010037 [arXiv:2112.02286 [astro-ph.IM]].
- [73] S. Hoof, J. Jaeckel and G. Lucente, [arXiv:2407.04772 [hep-ph]].
- [74] M. Giannotti, I. Irastorza, J. Redondo and A. Ringwald, JCAP **05**, 057 (2016) doi:10.1088/1475-7516/2016/05/057 [arXiv:1512.08108 [astro-ph.HE]].
- [75] M. Giannotti, I. G. Irastorza, J. Redondo, A. Ringwald and K. Saikawa, JCAP **10**, 010 (2017) doi:10.1088/1475-7516/2017/10/010 [arXiv:1708.02111 [hep-ph]].
- [76] L. Di Luzio, M. Fedele, M. Giannotti, F. Mescia and E. Nardi, JCAP **02**, no.02, 035 (2022) doi:10.1088/1475-7516/2022/02/035 [arXiv:2109.10368 [hep-ph]].
- [77] S. Hoof, J. Jaeckel and L. J. Thormaehlen, JCAP **10**, 024 (2023) doi:10.1088/1475-7516/2023/10/024 [arXiv:2306.00077 [hep-ph]].
- [78] J. Jaeckel and L. J. Thormaehlen, Phys. Rev. D **100**, no.12, 123020 (2019) doi:10.1103/PhysRevD.100.123020 [arXiv:1908.10878 [astro-ph.SR]].
- [79] S. Hoof and J. Jaeckel, Phys. Rev. D **96**, no.11, 115016 (2017) doi:10.1103/PhysRevD.96.115016 [arXiv:1709.01090 [hep-ph]].
- [80] M. Xiao, P. Carenza, M. Giannotti, A. Mirizzi, K. M. Perez, O. Straniero and B. W. Grefenstette, Phys. Rev. D **106**, no.12, 123019 (2022) doi:10.1103/PhysRevD.106.123019 [arXiv:2204.03121 [astro-ph.HE]].

- [81] M. Xiao, K. M. Perez, M. Giannotti, O. Straniero, A. Mirizzi, B. W. Grefenstette, B. M. Roach and M. Nynka, *Phys. Rev. Lett.* **126**, no.3, 031101 (2021) doi:10.1103/PhysRevLett.126.031101 [arXiv:2009.09059 [astro-ph.HE]].
- [82] F. Arias-Aragón, M. Giannotti, G. G. di Cortona and F. Mescia, [arXiv:2411.19327 [hep-ph]].
- [83] P. Carenza, G. Co', T. Fischer, M. Giannotti, A. Mirizzi and T. Rauscher, doi:10.3204/PUBDB-2018-02580 [arXiv:1808.04101 [hep-ph]].
- [84] A. Lella, P. Carenza, G. Lucente, M. Giannotti and A. Mirizzi, *Phys. Rev. D* **107**, no.10, 103017 (2023) doi:10.1103/PhysRevD.107.103017 [arXiv:2211.13760 [hep-ph]].
- [85] A. Lella, P. Carenza, G. Co', G. Lucente, M. Giannotti, A. Mirizzi and T. Rauscher, *Phys. Rev. D* **109**, no.2, 023001 (2024) doi:10.1103/PhysRevD.109.023001 [arXiv:2306.01048 [hep-ph]].
- [86] F. Calore, P. Carenza, C. Eckner, M. Giannotti, G. Lucente, A. Mirizzi and F. Sivo, *Phys. Rev. D* **109**, no.4, 043010 (2024) doi:10.1103/PhysRevD.109.043010 [arXiv:2306.03925 [astro-ph.HE]].
- [87] P. Carenza, G. Co, M. Giannotti, A. Lella, G. Lucente, A. Mirizzi and T. Rauscher, *Phys. Rev. C* **109**, no.1, 015501 (2024) doi:10.1103/PhysRevC.109.015501 [arXiv:2306.17055 [hep-ph]].
- [88] A. Lella, F. Calore, P. Carenza, C. Eckner, M. Giannotti, G. Lucente and A. Mirizzi, *JCAP* **11**, 009 (2024) doi:10.1088/1475-7516/2024/11/009 [arXiv:2405.02395 [hep-ph]].