

HAWC response to atmospheric electricity activity

Alejandro Lara*

Instituto de Geofísica, Universidad Nacional Autónoma de México

E-mail: alara@igeofisica.unam.mx

Graciela Binimelis de Raga

Centro de Ciencias de la Atmósfera, Universidad Nacional Autónoma de México

Olivia Enríquez-Rivera

Instituto de Geofísica, Universidad Nacional Autónoma de México

for the HAWC collaboration

The HAWC Gamma Ray observatory consists of 300 water Cherenkov detectors (WCD) instrumented with four photo multipliers tubes (PMT) per WCD. HAWC is located between two of the highest mountains in Mexico. The high altitude (4100 m asl), the relatively short distance to the Gulf of Mexico (~100 km), the large detecting area (22 000 m²) and its high sensitivity, make HAWC a good instrument to explore the acceleration of particles due to the electric fields existing inside storm clouds. In particular, the scaler system of HAWC records the output of each one of the 1200 PMTs as well as the 2, 3, and 4-fold multiplicities (logic AND in a time window of 30 ns) of each WCD with a sampling rate of 40 Hz. Using the scaler data, we have identified 20 enhancements of the observed rate during periods when storm clouds were over HAWC but without cloud-earth discharges. These enhancements can be produced by electrons with energy of tens of MeV, accelerated by the electric fields of tens of kV/m measured at the site during the storm periods. In this work, we present the recorded data, the method of analysis and our preliminary conclusions on the electron acceleration by the electric fields inside the clouds.

35th International Cosmic Ray Conference — ICRC2017

10–20 July, 2017

Bexco, Busan, Korea

*Speaker.

1. Introduction

Particle acceleration up to high energies inside the Earth's atmosphere has been observed by satellite gamma ray detectors during terrestrial gamma ray flashes [1, 2, 3, 4]. At ground level, high altitude cosmic ray detectors have reported ground enhancements during thunderstorms [5, 6, 7, 8].

The development of large electric fields during thunderstorms (up to 200 kV/m [9]) accelerates charged particles. Electrons may gain energies up to tens of MeV [7]. In this work we report enhancements of the count rates observed by the High Altitude Water Cherenkov Observatory (HAWC) which might be related to the atmospheric electric field.

HAWC is an air shower detector located at 4,100 m a.s.l, N 18°59'48", W 97°18'34". Built on the slope of Sierra Negra, Puebla in Mexico it consists of 300 water Cherenkov detectors 7.3 m diameter and 4.5 m deep. Each tank is filled with filtered water and the total detector comprises an extension of 22,000 m².

HAWC is operating on one of the highest mountains in Mexico. HAWC's high altitude together with the proximity of the Gulf of Mexico make the array an excellent laboratory to study the high energetic processes during thunderstorms. In section 2 we discuss the weather on the HAWC site in more detail.

In section 3 we describe HAWC scaler systems which are able to detect low energy particles. In Sec. 4 we present the enhancements of the scaler count rates observed by HAWC due to the presence of strong electric fields. Finally our discussion is presented in Sec. 5

2. The weather at the HAWC site

Southern and Central Mexico is located in the tropics, with ample humidity during most of the year and characterized by a 6-month rainy season. HAWC in particular, is located in a region with frequent presence of clouds formed by forced orographic lifting or due to atmospheric convective instability. The latter mechanism often leads to cumulus clouds responsible for the development of precipitation and are also responsible for charge separation within the cloud, due to collisions between hydrometeors at the different temperature ranges observed. The development of poles of positive and negative charge within the cloud give rise to an electric field that can reach breakdown point and result in a lightning discharge, within the cloud or from the cloud to ground.

Several ground-based networks have been developed to monitor cloud-to-ground lightning continuously to assess the risk to the population. One such global network is the World Wide Lightning Location Network (WWLN), documented by Dowden et al (2008). The first studies of combined precipitation and lightning over Mexico were carried out in 2010 and revealed the regions of the country where most of the cloud-to-ground lightning is observed (Kucienska et al, 2010). Furthermore, Raga et al (2014) showed that Mexico is particularly vulnerable, with a large number of deaths per year.

While HAWC is not located in the region of highest incidence of lightning in Mexico, its location on the Sierra Negra ensures that it will be affected by electrically-charged clouds and lightning for about 6 months of the year. The electric fields associated with the convective clouds and the proximity with intra-cloud and cloud-to-ground lightning provide a unique setting to study the effect of these phenomena on the measurements made by HAWC. In this work, we carried

41 out measurements of the ambient variables using a weather station placed at the HAWC site. To
 42 measure the electric field, we used an electric field mill Boltek-100 installed at the eastern side of
 43 the array. Plots of these variables are shown in section 4.




Figure 1: Average spatial distribution of cloud-to-ground lightning density in flashes per square kilometer per year for the period 2006-2012, adapted from Raga et al (2014). The white star mark the location of HAWC.

44 3. HAWC scaler systems

45 HAWC data are collected by two data acquisition systems (DAQs). The main DAQ measures
 46 arrival times and time over thresholds of PMT pulses and allows for the reconstruction of the air
 47 shower arrival direction and energy of the primary particle. The electronics are based on time
 48 to digital converters (TDC). The main DAQ also has a TDC scaler system which counts the hits
 49 inside a time window of 30 ns of each PMT and the coincidences of 2, 3 and 4 PMTs in each
 50 water Cherenkov detector. These coincidences are called multiplicity 2, 3 and 4, respectively.
 51 The secondary DAQ consists of a counting system that registers each time the PMT is hit by
 52 $> 1/4$ photoelectron charge and we call it hardware (HW) scaler system. This simpler system
 53 together with the TDC scalers allows one to measure particles below the energies of reconstructable
 54 showers.

55 4. Count rate enhancements

56 We have noted that the HAWC scaler systems responds to the atmospheric electricity at least
 57 in four ways: i) when the electric field is positive or weak negative the count rate does not suffer
 58 any change as seen in Figure 2 where we have plotted the TDC scaler rates during the negative
 59 electric field enhancement observed on Nov 22, 2014. All scaler rates are in percentage taking
 60 as reference, i.e. 100% , the mean scaler rate calculated one hour before the event started. The
 61 electric field is shown in black solid line. As an eye aid, the equivalent zero electric field is plotted




Figure 2: Mean count rate of the TDC scaler Multiplicities: 2 (blue), 3 (cyan) and 4 (green); and rates of the 8'' (orange) and 10'' (purple) PMTs, during November 22, 2014 when a moderate negative electric field was observed (black curve).

62 as a horizontal dashed line. ii) when the storm is very close to the array with large amount of
 63 discharges, the system behavior is unstable and the system restarts frequently. There are scaler
 64 enhancements but they may be due to the discharges and/or the electric field or electromagnetic
 65 noise. For example, Figure 3 shows the TDC scaler rates during November 8, 2014, an active day
 66 in terms of atmospheric electricity. A thunderstorm took place during this period as seen by the red
 67 square symbols and purple triangles representing the cloud to ground and inter-cloud discharges,
 68 respectively. There were nearby discharges as shown by the rapid variations of the field strength
 69 (it is important to note that the electric field detector gets saturated when the electric field is larger
 70 than ± 40 kV/m).




Figure 3: TDC scaler rates and electric field measured during Nov 8, 2014. The color code is similar as Fig. 2. The distance of the reported cloud to ground (red squares) and inter-cloud (purple triangles) discharges are also plotted.

71 There are some events where the atmospheric electric activity is not so strong and therefore,
 72 the response of the scaler system is somehow “well behaved.” In those cases we can distinguish:
 73 iii) a fast response of the scaler system associated to the discharges and iv) a slow response to
 74 strong negative electric fields. These responses are depicted in Figure 4 where we have plotted,

75 with colored dots, the count rate (in percentage) of each channel of the HW scaler system during
 76 September 18, 2015. The electric field shows two rapid changes around 00:45, associated with
 77 discharges. The squares and triangles indicate that a storm took place. There are sharp scaler
 78 enhancements associated with the closest discharges between 00:35 and 00:50 UT.




Figure 4: Mean count rate of the HW scaler system, each available channel is plotted in colored dots. The distance of the reported cloud to ground (red squares) and inter-cloud (purple triangles) discharges are also plotted.

79 In this work we focus on the slow response of the scaler system due to the presence of strong
 80 negative electric field. A clear example of this slow response was observed during May 26, 2015
 81 and is depicted in Figures 5 and 6 for TDC and HW scalers, respectively. We have selected this
 82 event due to the fact that there is no saturation of the electric field measurements.




Figure 5: Mean count rate of the TDC scaler Multiplicities during May 26, 2015. The color code is the same as in Figure 2.

83 The environment parameters during May 26, 2015 are presented in Figure 7. From top to
 84 bottom, we plotted the electric field, pressure, temperature, humidity, rain fall and solar irradiance.
 85 The latter two are displayed in order to show the presence of clouds at the site during the scaler
 86 enhancements. In particular, it is well known that the count rate of cosmic ray detectors has an
 87 inverse dependence on the ambient pressure. This figure shows that the count rate enhancements
 88 during the events are not related to pressure changes.

POS (ICRC2017) 080




Figure 6: Mean count rate of the HW scaler system during May 26, 2015. The color code is the same as in Figure 4.




Figure 7: Environment variables at the HAWC site during May 26, 2015. From top to bottom are: Electric field, ambient pressure, temperature (left panels), humidity, rainfall and solar irradiance (right panels).

89 Figure 8 shows the scatter plot of the TDC scaler enhancement as a function of the strength
 90 of the electric field, during the May 26 event. Unfortunately the electric field data are stored with
 91 poor time resolution (~ 1.5 minutes), limiting the statistics available for correlations. In order to
 92 show the tendency of the correlation, we fit a second degree polynomial to each mean multiplicity
 93 and mean PMT rates. The correlation coefficients are shown in the plot as reference. The scatter is
 94 high but one can see that both 10" PMT and multiplicity 2 rates are more affected by the electric
 95 field enhancement. The 8" PMTs, multiplicity 3 and finally multiplicity 4 are less affected. If the
 96 rate enhancements are being produced by the acceleration of charged particles in the electric field,
 97 this correlation would indicate more low-energy and few high-energy particles in the enhancement.
 98 Finally, the small scatter plots in Figure 8 emphasize the lack of correlation between the scaler rates
 99 and the Pressure/Temperature measured at the site.

100 **5. Discussion**

101 In this work, we presented examples of the HAWC scaler system response to the atmospheric




Figure 8: Scatter plot of the TDC scale Multiplicity 2 (green), 3 (blue) 4 (magenta); and 8'' (black) and 10'' PMT rates as a function of the electric field during May 26, 2015. The correlation coefficients are shown next to each Multiplicity and PMT set. Upper panels: scatter plots of the scaler rates vs Pressure (left) and Temperature (right)

102 electricity activity. We showed an example of the rapid response of the HW scaler system due to a
 103 close lightning activity (Figure 4). However, in this work we focus in the scaler system response to
 104 the negative electric field.

105 In particular we presented an example in which the electric field was not saturated and was
 106 observed by the two scaler systems. The correlated enhancement of all the available PMTs of the
 107 array seen by the HW scaler system (Figs. 6) shows that the enhancement embraces the entire
 108 array with similar response at time scales of seconds. The preliminary correlation analysis be-
 109 tween the negative electric field and TDC count rates shows the high relationship between these
 110 variables (Figures 8). Furthermore, the absence of correlation between the scaler count rates and
 111 atmospheric variables such as pressure or temperature supports a possible scenario where the scaler
 112 rate enhancements might be produced by particle acceleration due to the electric field of clouds ob-
 113 served by HAWC. If our hypothesis is correct, the enhancements of all multiplicities of the TDC
 114 scaler system (Fig. 5) will allow us to determine the energy of the incident particles.

115 We have shown that HAWC can be a good instrument to study the acceleration of particles
 116 by the atmospheric electricity. It is necessary to perform a detailed analysis and simulations to
 117 quantify our observations, as well as rule out instrumental effects in the photomultipliers (such as
 118 inductive charging) that could be producing the observed rate enhancements. This analysis will be
 119 published elsewhere.

120 Acknowledgments

121 We acknowledge the support from: the US National Science Foundation (NSF); the US De-
 122 partment of Energy Office of High-Energy Physics; the Laboratory Directed Research and De-
 123 velopment (LDRD) program of Los Alamos National Laboratory; Consejo Nacional de Ciencia
 124 y Tecnología (CONACyT), México (grants 271051, 232656, 260378, 179588, 239762, 254964,
 125 271737, 258865, 243290, 132197), Laboratorio Nacional HAWC de rayos gamma; L'OREAL Fel-
 126 lowship for Women in Science 2014; Red HAWC, México; DGAPA-UNAM (grants IG100317,
 127 IN111315, IN111716-3, IA102715, 109916, IA102917); VIEP-BUAP; PIFI 2012, 2013, PRO-
 128 FICIE 2014, 2015; the University of Wisconsin Alumni Research Foundation; the Institute of

129 Geophysics, Planetary Physics, and Signatures at Los Alamos National Laboratory; Polish Sci-
130 ence Centre grant DEC-2014/13/B/ST9/945; Coordinaci3n Cient3fica de la
131 Universidad Michoacana. Thanks to Luciano D3az and Eduardo Murrieta for technical support.

132 References

- 133 [1] G. Fishman, P. Bhat, R. Mallozzi, J. Horack, T. Koshut, C. Kouveliotou, G. Pendleton, C. Meegan, R.
134 Wilson, W. Paciesas, et al., *Science* 264, 1313 (1994).
- 135 [2] D. M. Smith, L. I. Lopez, R. P. Lin, and C. P. Barrington-Leigh, *Science* 307, 1085 (2005)
- 136 [3] M. Marisaldi, F. Fuschino, C. Labanti, M. Galli, F. Longo, E. Del Monte, G. Barbiellini, M. Tavani,
137 A. Giuliani, E. Moretti, et al., *J. Geophys. Res.* 115, A00E13 (2010)
- 138 [4] M. S. Briggs, G. J. Fishman, V. Connaughton, P. N. Bhat, W. S. Paciesas, R. D. Preece, C.
139 Wilson-Hodge, V. L. Chaplin, R. M. Kippen, A. von Kienlin, et al., *J. Geophys. Res.* 115, A07323
140 (2010) [11]
- 141 [5] V. V. Alexeenko, N.S. Khaerdinov, A.S. Lidvansky and V.B.Petkov, Transient variations of secondary
142 cosmic rays due to atmospheric electric field and evidence for pre-lightning particle acceleration,
143 *Phys. Lett. A*, 301 (2002) [12]
- 144 [6] S. Vernetto for EAS-TOP Collaboration, The EAS counting rate during thunderstorms. Proc. of 27th
145 ICRC,Copernicus Gesellschaft, Hamburg, Germany,August 7-15, 10 (2001) [17]
- 146 [7] A. Chilingarian, B. Mailyan and L. Vanyan, Recovering of the energy spectra of electrons and gamma
147 rays coming from the thunderclouds, *Atmos. Res.*, 114-115 (2012)
148 [18]
- 149 [8] H. Tsuchiya, K. Hibino and K. Kawata et al., Observation of thundercloud-related gamma rays and
150 neutrons in Tibet, *Phys. Rev. D*, 85 (2012)
- 151 [9] M. Stolzenburg, T. C. Marshall, W. D. Rust, E. Bruning, D. R. MacGorman and T. Hamlin, Electric
152 field values observed near lightning flash initiations, *Geophys. Res. Lett.* 34 (2007) L04804.
- 153 [10] Boccippio, Dennis, J., Kenneth L. Cummins, Hugh J. Christian, and Steven J. Goodman, 2001:
154 Combined Satellite- and Surface-Based Estimation of the Intracloud to Cloud-to-Ground Lightning
155 Ratio over the Continental United States. *Mon. Wea. Rev.*, 129, 1088-1122.
- 156 [11] Christian, H. J., Blakeslee R. J., Boccippio D. J., Boeck W. L., Buechler D. E., Driscoll K. T.,
157 Goodman S. J., Hall J. M., Koshak W. J., Mach D. M., and Stewart M. F., 2003, Global frequency and
158 distribution of lightning as observed from space by the Optical Transient Detector. *J. Geophys. Res.*,
159 108(D1), 4005.
- 160 [12] Dowden, R. and 30 more authors, 2008: World-Wide Lightning Location using VLF propagation in
161 the Earth-Ionosphere waveguide. *IEEE Antennas and Propagation Mag.*, 50, 40-60.
- 162 [13] Kucienska, B., G. B. Raga, and O. Rodriguez, 2010: Cloud-to-ground lightning over Mexico and
163 adjacent oceanic regions: a preliminary climatology using the WWLLN dataset. *Ann. Geophys.*, 28,
164 2047-2057.
- 165 [14] Raga, G.B., M.G. de la Parra and B. Kucienska, 2014: Deaths by lightning in Mexico (1979-2011):
166 Threat or vulnerability? *Wea. Climate Soc*, 6, 434-444.